

REVIEW ARTICLE

# Emerging endocrine disruptors: perfluoroalkylated substances

Allan Astrup Jensen\* and Henrik Leffers†

\*FORCE Technology, Broendby, Denmark, and †Department of Growth and Reproduction, Rigshospitalet, Copenhagen, Denmark

## Keywords:

endocrine disruption, fluorotelomers, perfluoroalkyl substances, perfluorooctane sulfonic acid, perfluorooctanoic acid, polyfluorinated chemicals

## Correspondence:

Allan Astrup Jensen, FORCE Technology, 345 Park Alle, DK-2605 Broendby, Denmark.  
E-mail: aaj@force.dk

Received 24 October 2007; revised 19 December 2007; accepted 8 January 2008

doi:10.1111/j.1365-2605.2008.00870.x

## Summary

In recent years, polyfluorinated chemicals (PFCs) have increasingly been used as surfactants in various industry- and consumer products, because of their unique properties as repellents of dirt, water and oils. The most well-known PFCs are perfluorooctane sulfonic acid (PFOS), perfluorooctanoic acid (PFOA) and their derivatives belonging to the group of perfluoroalkylated substances. The PFCs are very persistent in the environment, and some of them have been discovered as global pollutants of air, water, soil and wildlife and even found in remote polar areas. Bioaccumulation occurs also in humans, and everybody in our society has traces of these PFCs in their blood and internal organs such as the liver, kidneys, spleen, gall bladder and testes. In the blood, PFOS and PFOA are bound to serum proteins. The acute toxicity of the polyfluorinated substances is moderate but some substances can induce peroxisome proliferation in rat livers and may change the fluidity of cell membranes. Some of these PFCs, such as PFOS and PFOA, are potential developmental toxicants and are suspected endocrine disruptors with effects on sex hormone levels resulting in lower testosterone levels and higher oestradiol level. Other PFCs have oestrogenic effects in cell cultures. The industrial production of PFOS and its derivatives stopped in 2000, and the European Union has banned most uses from the summer of 2008. However, hundreds of related chemicals: homologues with shorter or longer alkyl chain, PFOA and telomers, which potentially may degrade to perfluoroalkanoic (carboxylic) acids, are not regulated.

## Introduction


Polyfluorinated chemicals (PFCs) are synthetic chemical substances that have very unique properties, such as high stabilities and extremely low surface tensions. Most PFCs are insoluble in both water and organic solvents, and they repel dirt, water and oils. Although these chemicals are rather expensive several hundreds different of them are increasingly used as surfactants in various industry- and common consumer products, including 'nanoproducts'. Important are uses in paints and for impregnation of textiles, clothes, footwear, furniture and carpets. Other uses are in lubricants, waxes for floors and cars, and in fire-fighting foam for oil fires at airports, harbours, oil platforms and oil refineries. Common trade names for commercial products are: Baygard (Bayer AG, Leverkusen, Germany), Scotchgard (3M, St. Paul, MN, USA), Gore-Tex (W. L. Gore & Associates Inc., Newark, DE, USA), Zonyl (E. I. du Pont de Nemours and Company, Wilmington,

DE, USA) and Stainmaster (INVISTA, Wichita, KS, USA) (Hekster *et al.*, 2003; Poulsen *et al.*, 2005; OECD, 2006).

The perfluoroalkylated substances (PFAS), such as perfluorooctane sulfonic acid (PFOS) and perfluorooctanoic acid (PFOA), are most persistent in the environment and have been discovered as global pollutants of air, water and soil, and even found in remote polar areas (Giesy & Kannan, 2002). PFOS is most abundant and have been determined in blood and liver samples from various aquatic mammals (seals, otters, sea lions, dolphins, polar bears and minks) and birds, fish and humans. The highest levels of PFOS have been found in polar bears from the Arctic (Bossi *et al.*, 2005; Smithwick *et al.*, 2005a,b, 2006; Betts, 2007).

## Chemistry and properties

The most well-known perfluoroalkyl substances are PFOS, PFOA and their derivatives. The chemical structures for some compounds are shown in Fig. 1.



**Figure 1** Chemical structures of selected polyfluorinated chemicals.

PFOA, PFOS and their salts are water-soluble and may be spread globally by sea water currents (Prevedouros *et al.*, 2006).

The substances most used in practice are more complex derivatives of these basic chemical structures, e.g. sulfonamides and fluorotelomers (Fig. 1).

These complex derivatives may be degraded to either PFOS, PFOA or perfluoroalkanoic acids with shorter or longer perfluorinated alkyl chain; the perfluorinated 'tail' is not degraded in the body or in the environment (Dinglasan *et al.*, 2004; Lau *et al.*, 2007).

The complex derivatives are insoluble in water and lipids but somewhat volatile and may be transported over long distances by air from temperate to arctic areas (Wallington *et al.*, 2006). The use of fluorotelomer derivatives in fire-fighting foam has an especially high potential for environmental release and human exposure to PFCs (Moody & Field, 2000; Moody *et al.*, 2002).

### Human exposure

Lipophilic persistent organic pollutants (POPs), such as DDT, polychlorinated biphenyls (PCB), dioxins and brominated diphenyl ethers, accumulate in fatty tissue, and the main route of exposure for the general population is from food intake (UNEP, 2001). However, PFCs are both lipophobic and hydrophobic and, therefore, they will not accumulate in lipids but mainly accumulate in blood, liver and kidneys, which are not so common food ingredients. A study from Canada has, however, reported PFAS in fast food composites (Tittlemier *et al.*, 2006).

The most publicly well-known occurrence of PFCs is probably as impurity in the non-stick surface layer of Teflon (E. I. du Pont de Nemours and Company) treated cookware, such as frying pans, and as greaseproof additive in pop corn microwave bags, from which the chemicals can be released gradually during cooking and leak into the food (Sinclair *et al.*, 2007).

The major human exposure is probably from PFCs used as surfactants for impregnation of consumer goods, such as textiles, foot wear, furniture and carpets, which then releases PFCs to the indoor air and contaminate indoor dust, which then is inhaled by humans. Indoor air contains in general 25–100 times more PFAS than outdoor air with maximum levels [e.g. 8 : 2 fluorotelomer alcohol (FTOH)] of  $28 \mu\text{g}/\text{m}^3$ , and house dust can be very contaminated with up to 75 ppm *N*-ethyl perfluorooctane sulfonamidoethanol (EtFOSE), 16 ppm 8 : 2 FTOH, 5 ppm PFOS and 3.7 ppm PFOA (Shoeib *et al.*, 2004, 2007). Moreover, children, who may be more in contact with considerable amounts of house dusts during play on the floor, will collect such dusts on the fingers, which subsequently can be ingested. Relative to body weight children have a 5–10 times larger intake of PFCs indoor than adults (Shoeib *et al.*, 2005).

### Human levels and half-lives

The PFAS are readily absorbed and bind to proteins in blood serum and accumulate mainly in organs such as liver, kidney and spleen, but also in testicles and brain (Vanden Heuvel *et al.*, 1992; Austin *et al.*, 2003; Jones *et al.*, 2003). The renal clearances of PFOA and PFOS is almost negligible in humans, contrary to a large active excretion in experimental animals (Harada *et al.*, 2005). The half-lives of different polyfluorinated compounds vary but are generally very long. The half-lives in blood serum have been estimated to be 3.8, 5.4 and 8.5 years for PFOA, PFOS and perfluorohexane sulfonamide, respectively (Olsen *et al.*, 2007) – and in the organs, the half-lives are probably even longer. Levels in serum are, in general, 2–3 times higher than in whole blood (Taniyasu *et al.*, 2003; Ehresman *et al.*, 2007).

Today PFAS, such as PFOA and PFOS, are present in the blood, liver and kidneys of everybody in our society (Kannan *et al.*, 2004). In utero exposure to PFOS and PFOA is, therefore, also ubiquitous as they are transferred

to the foetus through the placenta and later also to the babies through the milk (Apelberg *et al.*, 2007a; HENDERSON & Smith, 2007).

Levels in blood are considered a sufficient sensitive, precise and accurate biomarker of human exposure to PFOS (Butenhoff *et al.*, 2006). Levels can be high in blood serum of the general population, as levels of more than 1500 ng PFOS/mL have been observed in American blood donors (Olsen *et al.*, 2003a). In general, average levels are, however, much lower around 20–30 ng PFOS/mL, and the levels of PFOA and other perfluorocarboxylic acids even lower. There are also geographical variations as illustrated in Fig. 2.

Today the levels are so high that exposure to polyfluorinated substances measured on basis of whole blood may be the highest human exposure to exogenous chemicals, exceeding that of more well-known environmental contaminants such as DDE (persistent and bioaccumulative metabolite of the pesticide DDT), PCBs, brominated flame retardants and even phthalates. Results from a study in the Norwegian and Russian Arctic are shown in Fig. 3.

Levels of polyfluorinated compounds in maternal blood are in general 2–3 times the levels in cord blood (Apelberg *et al.*, 2007a; Fei *et al.*, 2007; Inoue *et al.*, 2004; Tittlemier *et al.*, 2004; Midasch *et al.*, 2007). Levels in semen were 10 times lower than in blood serum (Guruge *et al.*, 2005).


## Toxicology

There is limited information available on the toxicology of most PFCs (Lau *et al.*, 2007), and it will probably last several years before there is sufficient knowledge to assess the full consequences of the human exposure to these chemicals.


The acute toxicity of PFAS is moderate but increases with the chain length. The oral rat LD<sub>50</sub> for PFOS is 251 mg/kg (US EPA, 2000), while the oral LD<sub>50</sub> for PFOA is between 430 and 1800 mg/kg (Kennedy *et al.*, 2004).

The primary target organ is the liver, where they cause an increased weight and hepatocytic hypertrophy (Seacat *et al.*, 2003; Kennedy *et al.*, 2004). In rats, some of these substances induce lower serum glucose and cholesterol levels and increases the  $\beta$ -oxidation of fatty acids (Sohleinius *et al.*, 1995; Seacat *et al.*, 2003). Moreover, PFAS binds to proteins in cell membranes and change the membrane properties (fluidity), which can lead to aberrant signalling from surface receptors (Hu *et al.*, 2003).

Perfluorooctane sulfonic acid exposure (10 mg/kg bw for 2 weeks) of female rats affects the oestrous cyclicity and increases serum corticosterone level, while it decreases serum leptin concentration and norepinephrine concentration in the hypothalamus (Austin *et al.*, 2003). In utero exposure to PFOS ( $\geq 0.8$  mg/kg bw/day) leads in



**Figure 2** Typical average concentrations of perfluorooctane sulfonic acid and perfluorooctanoic acid in blood (serum/plasma) from various countries (Hansen *et al.*, 2001; Olsen *et al.*, 2003a,b, 2004, 2007; Corsolini & Kannan, 2004; Kannan *et al.*, 2004; Kubwabo *et al.*, 2004; Kuklennyik *et al.*, 2004; Tittlemier *et al.*, 2004; Guruge *et al.*, 2005; Calafat *et al.*, 2006; De Silva & Mabury, 2006; Odland *et al.*, 2006; Thomsen *et al.*, 2006; Fei *et al.*, 2007; Fromme *et al.*, 2007; Harada *et al.*, 2007; Kärrman *et al.*, 2007; Kato *et al.*, 2007).



**Figure 3** Persistent organic pollutants in blood plasma from pregnant women living in Norwegian and Russian Arctic (modified from Odland *et al.*, 2006).

mice and rats dose-dependently to a reduced litter size caused both by reduced implantation and foetal and neonatal death (Luebker *et al.*, 2005a; Lau *et al.*, 2006). Pups of dams of mice and rats exposed by oral gavage to PFOS ( $\geq 0.4$  mg/kg bw/day) showed a reduced body weight and a delay in development and neuromotor maturation (Luebker *et al.*, 2005b; Fuentes *et al.*, 2007), but there are no reports of increase in malformations (Lau *et al.*, 2007). Gestational PFOA exposure of mice is associated with altered mammary gland development in Dams and female offspring (White *et al.*, 2007).

Apelberg *et al.* (2007b) found that reductions of birth weight, ponderal index and head circumference were associated with PFOS and PFOA concentrations in blood among vaginal deliveries. A similar inverse association concerning maternal PFOA concentration and birth weight was found by Fei *et al.* (2007).

Perfluorooctane sulfonic acid, PFOA and other tested PFCs were not active in various mutagenicity test systems. However, PFOA exposed rodents developed an excess of Leydig cell adenomas (Cook *et al.*, 1992; Biegel *et al.*, 1995; Liu *et al.*, 1996). Moreover, PFOS and EtFOSE caused liver hepatocellular adenomas in female rats and thyroid follicular cell adenomas in male rodents (Thomford *et al.*, 2002). Because of these studies, the US EPA classifies PFOA as an animal carcinogen (US EPA, 2000). In accordance, a study from the work environment reports an increased incidence of urinary bladder cancer in workers with high exposure to perfluorooctane sulfonyl fluoride (Alexander *et al.*, 2003).

Thus, although PFAS are only moderately toxic, they do induce a number of adverse effects in experimental animals, which together point to potential problems with the currently very high human exposure levels.

### Possible mixture effects

As exposure to PFCs is ubiquitous to humans this exposure is added to all other exposures humans may experience. This raises the question of possible mixture effects, which could increase effects from other exposures. Mixture effects have already been shown in vitro and in vivo for oestrogenic compounds (Silva *et al.*, 2002; Tinwell & Ashby, 2004) and for anti-androgens (Birkhoj *et al.*, 2004; Metzdorff *et al.*, 2007), but there are very few studies of mixture effects of compounds with different modes of action.

For PFAS, there are several reports of such effects. Co-administration of PFOS and dioxin [2378-tetrachlorodibenzo-p-dioxin (TCDD)] resulted in an increased p450 A1A expression as compared to TCDD alone (Hu *et al.*, 2003). PFAS may also increase the carcinogenicity of other chemicals as the genotoxicity of cyclophosphamide in the micronucleus assay with hamster lung V79 cells increased many fold by simultaneous exposure to PFOS (Jernbro *et al.*, 2007).

Thus, the sensitivity to other chemicals can be increased by simultaneous exposure to PFCs.

### Endocrine disruption

Some PFCs have oestrogenic effects in cell cultures ('E-screen assay'; Soto *et al.*, 1995). For example, the fluorotelomer alcohols 6 : 2 FTOH and 8 : 2 FTOH induce MCF-7 breast cancer cell proliferation and up-regulates the oestrogen receptor, but PFOS, PFOA and perfluorononanoic acid had no oestrogenic effect in that test (Maras *et al.*, 2006; Vanparys *et al.*, 2006). Nevertheless, PFCs act as endocrine disruptors, because exposure of adult rats affects their endocrine system by decreasing the

testosterone level and increasing the oestradiol level (Biegel *et al.*, 1995; Shi *et al.*, 2007).

The effects on hormone levels in rodents are reflected in changes in the testis where exposure to perfluorooctanoate results in Leydig cell hyperplasia and eventually the development of Leydig cell adenomas (Biegel *et al.*, 1995). A study of testis effects in adult rats exposed to  $\geq 5$  mg perfluorododecanoic acid/kg bw daily for 2 weeks also showed a reduced gene expression of many genes involved in cholesterol transport and steroidogenesis and a reduced serum testosterone level (Shi *et al.*, 2007). Thus, it seems as if exposure to PFAS substances can severely affect proliferation and function of Leydig cells in the adult rat.

This is of considerable concern, because Leydig cell hyperplasia is common among infertile men (Holm *et al.*, 2003) who, as a group, also show lower testosterone levels than comparable normal controls (Andersson *et al.*, 2004). Reduced testis function has been linked to the testicular dysgenesis syndrome (TDS) (Skakkebaek *et al.*, 2001). The TDS hypothesis states that in utero exposure to endocrine disruptors can damage testis development and lead to reduced testis function in the adult, with symptoms ranging from a moderately reduced semen quality to testis cancer. The best animal model for human TDS consists of rats exposed to long-chain phthalates in a critical time window during development (Fisher *et al.*, 2003). The exposure results in testis dysgenesis with Leydig cell hyperplasia and clustering of the Leydig cells in the centre of the testis, resulting in reduced testosterone levels and compromised fertility in the adults (Sharpe, 2006; Hallmark *et al.*, 2007). The compromised Leydig cell function is reflected in a reduced expression of genes involved in cholesterol transport and steroidogenesis (Liu *et al.*, 2005). This has striking similarities to the reported effect of PFAS exposure; however, it seems as if PFAS compounds, in contrast to phthalates, can induce the effects in the adult.

## Conclusion

Polyfluorinated chemicals have increasing importance in industry and consumer products because of their unique properties. However, these chemicals are persistent and have become wide-spread in the environment and accumulate in wildlife and humans.

One of the most worrying effects of polyfluorinated substances is their effect on Leydig cells in the rat testis, where the reported effects have a strong resemblance to observations from the clinic of infertile men seeking help with assisted reproduction. Such effects (Leydig cell hyperplasia, lower testosterone levels) can also be induced by exposure to other ubiquitous chemicals such

as phthalates. As PFCs can induce these effects in the adult, it raises the question of what happens when a person with a moderately reduced testis function, maybe caused by TDS induced in utero, as an adult is exposed to chemicals that can elicit the same effect.

Butenhoff *et al.* (2004) have made a risk characterization of the general population exposure to PFOA concluding that the margin of safety was ranging between 1600 for liver weight increase and 8900 for Leydig cell adenomas. This assessment was based on information obtained from animal experiments. However, the facts that the half-life of the chemical in humans is much longer than in animals, and the renal clearance in humans is negligible, contrary to the active excretion in animals, makes human risk assessment based on animal experiments questionable for these chemicals.

The international awareness and concern is increasing. In 2000 the main producer, the 3M Company, voluntarily stopped the production of one of the chemicals (PFOS), and a ban of some fluorotelomers has been introduced in Canada. In Europe, the EU countries will ban PFOS and its derivatives from the summer of 2008. However, PFOS is only a small part of the problem. The family of PFCs consists of several hundreds other unrestricted chemicals. Because the exposure to polyfluorinated substances is so considerable, and uses seem to increase, there is an urgent need to resolve, what effect such exposure has on humans.

## References

- Alexander, B. J., Olsen, G. W., Burris, J. M., Mandel, J. H. & Mandel, J. S. (2003) Mortality of employees of a perfluorooctane sulfonyl fluoride manufacturing facility. *Occupational and Environmental Medicine* 60, 722–729.
- Andersson, A. M., Jorgensen, N., Frydelund-Larsen, L., Rajpert-De Meyts, E. & Skakkebaek, N. E. (2004) Impaired Leydig cell function in infertile men: a study of 357 idiopathic infertile men and 318 proven fertile controls. *Journal of Clinical Endocrinology and Metabolism* 89, 3161–3167.
- Apelberg, B. J., Goldman, L. R., Calafat, A. M., Herbstman, J. B., Kuklenyik, Z., Heidler, J., Needham, L. L., Halden, R. U. & Witter, F. R. (2007a) Determinants of fetal exposure to polyfluoroalkyl compounds in Baltimore, Maryland. *Environmental Science and Technology* 41, 3891–3897.
- Apelberg, B. J., Witter, F. R., Herbstman, J. B., Calafat, A. M., Halden, R. U., Heidler, J., Needham, L. L. & Goldman, L. R. (2007b) Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in relation to weight and size at birth. *Environmental Health Perspectives* 115, 1670–1676.
- Austin, M. E., Kasturi, B. S., Barber, M., Kannan, K., MohanKumar, P. S. & MohanKumar, S. M. J. (2003)

- Neuroendocrine effects of perfluorooctane sulfonate in rats. *Environmental Health Perspectives* 111, 1485–1489.
- Betts, K. S. (2007) Perfluoroalkyl acids. What is the evidence telling us? *Environmental Health Perspectives* 115, A250–A256.
- Biegel, L. B., Liu, R. C. M., Hurtt, M. E. & Cook, J. C. (1995) Effects of ammonium perfluorooctanoate on Leydig cell function: in vitro, in vivo and ex vivo studies. *Toxicology and Applied Pharmacology* 134, 18–25.
- Birkhoj, M., Nellemann, C., Jarfelt, K., Jacobsen, H., Andersen, H. R., Dalgaard, M. & Vinggaard, A. M. (2004) The combined antiandrogenic effects of five commonly used pesticides. *Toxicology and Applied Pharmacology* 201, 10–20.
- Bossi, R., Riget, F. F., Dietz, R., Sonne, C., Fauser, P., Dam, M. & Vorkamp, K. (2005) Preliminary screening of perfluorooctane sulfonate (PFOS) and other fluorochemicals in fish, birds and marine animals from Greenland and the Faroe Islands. *Environmental Pollution* 136, 323–329.
- Butenhoff, J. L., Gaylor, D. W., Moore, J. A., Olsen, G. W., Rodricks, J., Mandel, J. H. & Zobel, L. R. (2004) Characterization of risk for general population exposure to perfluorooctanoate. *Regulatory Toxicology and Pharmacology* 39, 363–380.
- Butenhoff, J. L., Olsen, G. W. & Pfahles-Hutchens, A. (2006) The applicability of biomonitoring data for perfluorooctane-sulfonate to the environmental public health continuum. *Environmental Health Perspectives* 114, 1776–1782.
- Calafat, A. M., Kuklenyik, Z., Caudill, S. P., Reidy, J. A. & Needham, L. L. (2006) Perfluorochemicals in pooled serum samples from United States residents in 2001 and 2002. *Environmental Science and Technology* 40, 2128–2134.
- Cook, J. C., Murray, S. M., Frame, R. S. & Hurtt, M. E. (1992) Induction of Leydig cell adenomas by ammonium perfluorooctanoate: a possible endocrine-related mechanism. *Toxicology and Applied Pharmacology* 1134, 209–217.
- Corsolini, S. & Kannan, K. (2004) Perfluorooctane sulfonate and related fluorochemicals in several organisms including humans from Italy. *Organohalogen Compounds* 66, 4079–4085.
- De Silva, A. O. & Mabury, S. A. (2006) Isomer distribution of perfluorocarboxylates in human blood: potential correlation to source. *Environmental Science and Technology* 40, 2903–2909.
- Dinglasan, M. J. A., Ye, Y., Edwards, E. A. & Mabury, S. A. (2004) Fluorotelomer alcohol biodegradation yields poly- and perfluorinated acids. *Environmental Science and Technology* 38, 2857–2864.
- Ehresman, D. J., Froelich, J. W., Olsen, G. W., Chang, S.-C. & Butenhoff, J. L. (2007) Comparison of whole blood, plasma, and serum matrices for the determination of perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA), and other fluorochemicals. *Environmental Research* 103, 176–184.
- Fei, C., McLaughlin, J. K., Tarone, R. E. & Olsen, J. (2007) Perfluorinated chemicals and fetal growth: a study within the Danish National Birth Cohort. *Environmental Health Perspectives* 115, 1677–1682.
- Fisher, J. S., Macpherson, S., Marchetti, N. & Sharpe, R. M. (2003) Human 'testicular dysgenesis syndrome': a possible model using in-utero exposure of the rat to dibutyl phthalate. *Human Reproduction* 18, 1383–1394.
- Fromme, H., Midasch, O., Twardella, D., Angerer, J., Boehmer, S. & Liebl, B. (2007) Occurrence of perfluorinated substances in an adult German population in Southern Bavaria. *International Archives of Occupational and Environmental Health* 80, 313–319.
- Fuentes, S., Colomina, M. T., Vicens, P., Franco-Pons, N. & Domingo, J. L. (2007) Concurrent exposure to perfluorooctane sulfonate and restraint stress during pregnancy in mice: effects on postnatal development and behavior of the offspring. *Toxicological Sciences* 98, 589–598.
- Giesy, J. P. & Kannan, K. (2002) Perfluorochemical surfactants in the environment. *Environmental Science and Technology* 36, 147A–152A.
- Guruge, K. S., Taniyasu, S., Yamashita, N., Wijeratna, S., Mohotti, K. M., Seneviratne, H. R., Kannan, K., Yamanaka, N. & Miyazaki, S. (2005) Perfluorinated organic compounds in human blood serum and seminal plasma: a study of urban and rural tea worker populations in Sri Lanka. *Journal of Environmental Monitoring* 7, 371–377.
- Hallmark, N., Walker, M., McKinnell, C., Mahood, I. K., Scott, H., Bayne, R. *et al.* (2007) Effects of monobutyl and di(n-butyl) phthalate in vitro on steroidogenesis and Leydig cell aggregation in fetal testis explants from the rat: comparison with effects in vivo in the fetal rat and neonatal marmoset and in vitro in the human. *Environmental Health Perspectives* 115, 390–396.
- Hansen, K. J., Clemen, L. A., Ellefson, M. E. & Johnson, H. O. (2001) Compound-specific, quantitative characterization of organic fluorochemicals in biological matrices. *Environmental Science and Technology* 35, 766–770.
- Harada, K., Inoue, K., Morikawa, A., Yoshinga, T., Saito, N. & Koizumi, A. (2005) Renal clearance of perfluorooctane sulfonate and perfluorooctanoate in humans and their species-specific excretion. *Environmental Research* 99, 253–261.
- Harada, K., Koizumi, A., Saito, N., Inoue, K., Yoshinaga, T., Date, C. *et al.* (2007) Historical and geographical aspects of the increasing perfluorooctanoate and perfluorooctane sulfonate contamination in human serum in Japan. *Chemosphere* 66, 293–301.
- Hekster, F. M., Laane, R. W. P. M. & de Voogt, P. (2003) Environmental and toxicity effects of perfluoroalkylated substances. *Reviews of Environmental Contamination and Toxicology* 179, 99–121.
- Henderson, W. & Smith, M. A. (2007) Perfluorooctanoic acid and perfluorononanoic acid in fetal and neonatal mice following in utero exposure to 8–2 fluorotelomer alcohol. *Toxicological Sciences* 95, 452–461.
- Holm, M., Rajpert-De Meyts, E., Andersson, A. M. & Skakkebaek, N. E. (2003) Leydig cell micronodules are a common finding in testicular biopsies from men with impaired

- spermatogenesis and are associated with decreased testosterone/LH ratio. *Journal of Pathology* 199, 378–386.
- Hu, W., Jones, P. D., De Coen, W., King, L., Fraker, P., Newsted, J. & Giesy, J. P. (2003) Alterations in cell membrane properties caused by perfluorinated compounds. *Comparative Biochemistry and Physiology. C, Comparative Pharmacology* 135, 77–88.
- Inoue, K., Okada, F., Ito, R., Kato, S., Sasaki, S., Nakajima, S. *et al.* (2004) Perfluorooctane sulfonate (PFOS) and related perfluorinated compounds in human maternal and cord blood samples: assessment of PFOS exposure in a susceptible population during pregnancy. *Environmental Health Perspectives* 112, 1204–1207.
- Jernbro, S., Rocha, P. S., Keiter, S., Skutlarek, D., Färber, H., Jones, P. D., Giesy, J. P., Hollert, H. & Engwall, M. (2007) Perfluorooctane sulfonate increases the genotoxicity of cyclophosphamide in the micronucleus assay with V79 cells. *Environmental Science and Pollution Research* 14, 85–87.
- Jones, P. D., Hu, W., De Coen, W., Newsted, J. & Giesy, J. P. (2003) Binding of perfluorinated fatty acids to serum proteins. *Environmental Toxicology Chemistry* 22, 2639–2649.
- Kannan, K., Corsolini, S., Falandysz, J., Fillman, G., Kumar, K. S., Loganathan, B. G. *et al.* (2004) Perfluorooctane sulfonate and related fluorochemicals in human blood from several countries. *Environmental Science and Technology* 38, 4489–4495.
- Kärman, A., Ericson, I., van Bavel, B., Darnerud, P. O., Aune, M., Glynn, A., Lignell, S. & Lindström, G. (2007) Exposure of perfluorinated chemicals through lactation: levels of matched human milk and serum and a temporal trend, 1996–2004, in Sweden. *Environmental Health Perspectives* 115, 226–230.
- Kato, K., Calafat, A. M., Reidy, J. A., Bryant, X. A., Nielsen, F., Weihe, P., Grandjean, P. & Needham, L. L. (2007) Serum concentrations of polyfluoroalkyl compounds in Faroe Island residents. *Organohalogen Compounds* 69, 161–164.
- Kennedy, G. L., Butenhoff, J. L., Olsen, G. W., O'Connor, J. C., Seacat, A. M., Perkins, R. G., Biegel, L. B., Murphy, S. R. & Farrar, D. G. (2004) The toxicology of perfluorooctanoate. *Critical Reviews in Toxicology* 34, 351–384.
- Kubwabo, C., Vais, N. & Benoit, F. M. (2004) A pilot study on the determination of perfluorooctanesulfonate and other perfluorinated compounds in blood of Canadians. *Journal of Environmental Monitoring* 6, 540–545.
- Kuklenyik, Z., Reich, J. A., Tully, J. S., Needham, L. L. & Calafat, A. M. (2004) Automated solid-phase extraction and measurement of perfluorinated organic acids and amides in human serum and milk. *Environmental Science and Technology* 38, 3698–3704.
- Lau, C., Thibodeaux, J. R., Hanson, R. G., Narotsky, M. G., Rogers, J. M., Lindstrom, A. B. & Strynar, M. J. (2006) Effects of perfluorooctanoic acid exposure during pregnancy in the mouse. *Toxicological Sciences* 90, 510–518.
- Lau, C., Anitole, K., Hodes, C., Lai, D., Pfahles-Hutchens, A. & Seed, J. (2007) Perfluoroalkyl acids: a review of monitoring and toxicological findings. *Toxicological Sciences* 99, 366–394.
- Liu, K., Lehmann, K. P., Sar, M., Young, S. S. & Gaido, K. W. (2005) Gene expression profiling following in utero exposure to phthalate esters reveals new gene targets in the etiology of testicular dysgenesis. *Biology of Reproduction* 73, 180–192.
- Liu, R. C. M., Hurtt, M. E., Cook, J. C. & Biegel, L. B. (1996) Effect of the peroxisome proliferator, ammonium perfluorooctanoate (C8), on hepatic aromatase activity in adult male Crl:CD BR (CD) rats. *Fundamental and Applied Toxicology* 30, 220–228.
- Luebker, D. J., York, R. G., Case, M. T., Hansen, K. J., Moore, J. A. & Butenhoff, J. L. (2005a) Neonatal mortality from in utero exposure to perfluorooctanesulfonate (PFOS) in Sprague-Dawley rats: dose-response, and biochemical and pharmacokinetic parameters. *Toxicology* 215, 149–169.
- Luebker, D. J., Case, M. T., York, R. G., Moore, J. A., Hansen, K. J. & Butenhoff, J. L. (2005b) Two-generation reproduction and cross-foster studies of perfluorooctanesulfonate (PFOS) in rats. *Toxicology* 215, 126–148.
- Maras, M., Vanparys, C., Muylle, F., Robbens, J., Berger, U., Barber, J. L., Blust, R. & De Coen, W. (2006) Estrogen-like properties of fluorotelomer alcohols as revealed by MCF-7 breast cancer cell proliferation. *Environmental Health Perspectives* 114, 100–105.
- Metzdorff, S. B., Dalgaard, M., Christiansen, S., Axelstad, M., Hass, U., Kiersgaard, M. K., Scholze, M., Kortenkamp, A. & Vinggaard, A. M. (2007) Dysgenesis and histological changes of genitals and perturbations of gene expression in male rats after in utero exposure to antiandrogen mixtures. *Toxicological Sciences* 98, 87–98.
- Midasch, O., Drexler, H., Hart, N., Beckmann, M. W. & Angerer, J. (2007) Transplacental exposure of neonates to perfluorooctanesulfonate and perfluorooctanoate: a pilot study. *International Archives of Occupational and Environmental Health* 80, 643–648.
- Moody, C. A. & Field, J. A. (2000) Perfluorinated surfactants and the environmental implication of their use in fire-fighting foams. *Environmental Science and Technology* 34, 3864–3870.
- Moody, C. A., Martin, J. W., Kwan, W. C., Muir, D. C. G. & Mabury, S. (2002) Monitoring perfluorinated surfactants in biota and surface water samples following an accidental release of fire-fighting foam into Etobicoke Creek. *Environmental Science and Technology* 36, 545–551.
- Odland, J. Ø., Sandanger, T. M. & Heimstad, E. S. (2006) Kartlegging av “nye” miljøgifter i human blodprøver fra Nord-Norge, Nord-Vest Russland og Sibir. SFT Rapport: 963/2006. Statens Forurensningstilsyn, Oslo. <http://www.sft.no/publikasjoner/overvaking/2184/ta2184.pdf>
- OECD (2006) Survey on Production and Use of PFOS, PFAS, PFOA, PFCA, Their Related Substances and Products/Mixtures Containing These Substances. Environment, Health

- and Safety Publications, Series on Risk Management No. 22; env/jm/mono(2006)36.
- Olsen, G. W., Hansen, K. J., Stevenson, L. A., Burris, J. M. & Mandel, J. H. (2003a) Human donor liver and serum concentrations of perfluorooctane sulfonate and other perfluorochemicals. *Environmental Science and Technology* 37, 888–891.
- Olsen, G. W., Church, T. R., Miller, J. P., Burris, J. M., Hansen, K. J., Lundberg, J. K. *et al.* (2003b) Perfluorooctane sulfonate and other fluorochemicals in the serum of American Red Cross adult blood donors. *Environmental Health Perspectives* 111, 1892–1901.
- Olsen, G. W., Church, T. R., Larson, E. B., van Belle, G., Lundberg, J. K., Hansen, K. J., Burris, J. M., Mandel, J. H. & Zobel, L. R. (2004) Serum concentrations of perfluorooctanesulfonate and other fluorochemicals in an elderly population from Seattle, Washington. *Chemosphere* 54, 1599–1611.
- Olsen, G. W., Burris, J. M., Ehresman, D. J., Froelich, J. W., Seacat, A. M., Butenhoff, J. L. & Zobel, L. R. (2007) Half-life of serum elimination of perfluorooctane sulfonate, perfluorohexane sulfonate and perfluorooctanoate in retired fluorochemicals production workers. *Environmental Health Perspectives* 115, 1298–1305.
- Poulsen, P. B., Jensen, A. A. & Wallström, E. (2005) More Environmentally Friendly Alternatives to PFOS-Compounds and PFOA. Environmental Project No. 1013. Danish Environmental Protection Agency, Copenhagen.
- Prevedouros, K., Cousins, I. T., Buck, R. C. & Korzeniowski, S. H. (2006) Sources, fate and transport of perfluorocarboxylates. *Environmental Science and Technology* 40, 32–44.
- Seacat, A. M., Thomford, P. J., Hansen, K. J., Clemen, L. A., Eldridge, S. R., Elcombe, C. R. & Butenhoff, J. L. (2003) Sub-chronic dietary toxicity of potassium perfluorooctane sulfonate in rats. *Toxicology* 183, 117–131.
- Sharpe, R. M. (2006) Pathways of endocrine disruption during male sexual differentiation and masculinization. *Best Practice and Research. Clinical Endocrinology and Metabolism* 20, 91–110.
- Shi, Z., Zhang, H., Liu, Y., Xu, M. & Dai, J. (2007) Alterations in gene expression and testosterone synthesis in the testes of male rats exposed to perfluorododecanoic acid. *Toxicological Sciences* 98, 206–215.
- Shoeib, M., Harner, T., Ikononou, M. & Kannan, K. (2004) Indoor and outdoor air concentrations and phase partitioning of perfluoroalkyl sulfonamides and polybrominated diphenyl ethers. *Environmental Science and Technology* 38, 1313–1320.
- Shoeib, M., Harner, T., Wilford, B. H., Jones, K. C. & Zhu, J. (2005) Perfluorinated sulfonamides in indoor and outdoor air and indoor dust: Occurrence, partitioning, and human exposure. *Environmental Science and Technology* 39, 6599–6606.
- Shoeib, M., Harner, T. & Zhu, J. (2007) Indoor air & dust concentration of fluorotelomer alcohols. *Organohalogen Compounds* 69, 146–149.
- Silva, E., Rajapakse, N. & Kortenkamp, A. (2002) Something from “nothing”—eight weak estrogenic chemicals combined at concentrations below NOECs produce significant mixture effects. *Environmental Science and Technology* 36, 1751–1756.
- Sinclair, E., Kim, S. K., Akinleye, H. B. & Kannan, K. (2007) Quantitation of gas-phase perfluoroalkyl surfactants and fluorotelomer alcohols released from non-stick cookware and microwave popcorn bags. *Environmental Science and Technology* 41, 1180–1185.
- Skakkebaek, N. E., Rajpert-De Meyts, E. & Main, K. M. (2001) Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Human Reproduction* 16, 972–978.
- Smithwick, M., Muir, D. C. G., Mabury, S. A., Solomon, K., Martin, J. W., Sonne, C., Born, E. W., Letcher, R. J. & Dietz, R. (2005a) Perfluoroalkyl contaminants in liver tissue from East Greenland polar bears (*Ursus maritimus*). *Environmental Toxicology Chemistry* 24, 981–986.
- Smithwick, M., Mabury, S. A., Solomon, K., Sonne, C., Martin, J. W., Born, E. W. *et al.* (2005b) Circumpolar study of perfluoroalkyl contaminants in polar bears (*Ursus maritimus*). *Environmental Science and Technology* 39, 5517–5523.
- Smithwick, M., Norstrom, R. J., Mabury, S. A., Solomon, K., Evans, T. J., Stirling, I., Taylor, M. K. & Muir, D. C. G. (2006) Temporal trends of perfluoroalkyl contaminants in polar bears (*Ursus maritimus*) from two locations in the North American Arctic, 1972–2002. *Environmental Science and Technology* 40, 1139–1143.
- Sohlenius, A.-K., Andersson, K., Olsson, J. & DePierre, J. W. (1995) Peroxisome proliferation and associated effects caused by perfluorooctanoic acid in vitamin A-deficient mice. *Chemico-Biological Interactions* 98, 45–50.
- Soto, A. M., Sonnenschein, C., Chung, K. L., Fernandez, M. F., Olea, N. & Serrano, F. O. (1995) The E-SCREEN assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environmental Health Perspectives* 103(Suppl 7), 113–122.
- Taniyasu, S., Kannan, K., Horii, Y., Hanari, N. & Yamashita, N. (2003) A survey of perfluorooctane sulfonate and related perfluorinated organic compounds in water, fish, birds, and humans from Japan. *Environmental Science and Technology* 37, 2634–2639.
- Thomford, P. J., Seacat, A. M. & Butenhoff, J. L. (2002) Terminal observations in Sprague-Dawley rats after lifetime dietary exposure to N-ethyl perfluorooctane sulfonamide ethanol. *Toxicologist* 66, 185.
- Thomsen, C., Kuklenyik, Z., Frøshaug, M. & Calafat, A. M. (2006) The ranges of polyfluorinated compounds in serum from a group of high consumers of fish from a contaminated lake in Norway. *Organohalogen Compounds* 68, 1701–1704.
- Tinwell, H. & Ashby, J. (2004) Sensitivity of the immature rat uterotrophic assay to mixtures of estrogens. *Environmental Health Perspectives* 112, 575–582.



- Tittlemier, S., Ryan, J. J. & Van Oostdam, J. (2004) Presence of anionic perfluorinated organic compounds in serum collected from northern Canadian populations. *Organohalogen Compounds* 66, 4009–4014.
- Tittlemier, S. A., Pepper, K. & Edwards, L. (2006) Concentrations of perfluorooctanesulfonamides in Canadian Total Diet Study composite food samples collected between 1992 and 2004. *Journal of Agricultural Food Chemistry* 54, 8385–8389.
- UNEP. (2001) Information Kit Stockholm 2001. Conference of Plenipotentiaries for the Stockholm Convention on Persistent Organic Pollutants, 22 to 23 May 2001 – Stockholm, Sweden. <http://www.pops.int/documents/press/infokit/en/Infokiten.htm>
- US EPA. (2000) Perfluorooctyl sulfonates; proposed significant new use rule. *Federal Register* 65, 62319–62333.
- Vanden Heuvel, J. P., Kuslikis, B. I. & Peterson, R. E. (1992) Covalent binding of perfluorinated fatty acids to proteins in the plasma, liver and testes of rats. *Chemico-Biological Interactions* 82, 317–328.
- Vanparys, C., Maras, M., Lenjou, M., Robbens, J., Van Bockstale, D., Blust, R. & De Coen, W. (2006) Flow cytometric cell cycle analysis allows for rapid screening of estrogenicity in MCF-7 breast cancer cells. *Toxicology in Vitro* 20, 1238–1248.
- Wallington, T. J., Hurley, M. D., Xia, J., Wuebbles, D. J., Sillman, S., Ito, A. *et al.* (2006) Formation of C7F15COOH (PFOA) and other perfluorocarboxylic acids during the atmospheric oxidation of 8 : 2 fluorotelomer alcohol. *Environmental Science and Technology* 40, 924–930.
- White, S. S., Calafat, A. M., Kuklenyik, Z., Villanueva, L. T., Zehr, R. D., Helfant, L. *et al.* (2007) Gestational PFOA exposure of mice is associated with altered mammary gland development in dams and female offspring. *Toxicological Sciences* 96, 133–144.

## Panel discussion

### S. Swan

You noticed a reduction in testosterone and altered Leydig cell differentiation in mammals after exposure to these chemicals. Did this have an effect on the anogenital distance (AGD)?

### A.A. Jensen

The AGD was not measured.

### O. Söder

The high level of PFAS in polar bears is worrying. What is the source of these chemicals, for example fish or environment? We should not expect the chemicals to be present in high concentrations in the environment of polar bears.

### A.A. Jensen

We do not know the origin of PFAS in polar bears. They are at the top in the food chain, but there is much less PFAS in the sea or in fish and seals. The main chemical in the bear is PFOS (perfluoro-octane sulphonic acid) but there are also some longer chain compounds and a few lower chain compounds. The extremely high PFOS levels may be linked to the biochemistry of the polar bears with high levels of vitamin A in the liver. The PFOS is predominantly present in the blood, kidney and liver, and some is transported to the testis and brain.