

Tracking the Pathways of Human Exposure to Perfluorocarboxylates

ROBIN VESTERGREN AND IAN T. COUSINS*

Department of Applied Environmental Science (ITM), Stockholm University,
Svante Arrhenius väg 8c, SE-10691 Stockholm, Sweden

Received January 22, 2009. Revised manuscript received May 14, 2009. Accepted June 15, 2009.

Recent analyses of perfluorooctanoate (PFOA) in human blood sera show that the background-exposed population in industrialized countries worldwide exhibits a narrow concentration range; arithmetic means of published studies range between 2 and 8 $\mu\text{g/L}$ PFOA, with the exception of a few outlier studies. The globally comparable human serum concentrations of PFOA and characteristic dominance of PFOA with respect to other perfluorocarboxylate (PFCA) homologues indicate that exposure pathways of humans differ from those of wildlife, where perfluorononanoate (PFNA) is often the dominant homologue. The observed correlations between perfluorooctane sulfonate (PFOS) and PFOA in human serum together with a simultaneous downward time trend of these compounds in human blood sera and blood spots from the year 2000 onward indicate a connection between historical perfluorooctanesulfonyl (POSF) production (phased out by the major manufacturer in 2000–2002) and exposure to both PFOS and PFOA. A comparison of estimated daily intakes to humans based on samples from exposure media (collected post 2000) indicates that food intake is the major contemporary exposure pathway for the background population, whereas drinking water exposure is dominant for populations near sources of contaminated drinking water. A one-compartment pharmacokinetic model used to back-calculate daily intakes from serum levels is shown to provide agreement within a factor of 1.5–5.5 of the daily intakes derived from exposure media, which provides further supporting evidence that dietary exposure is a major ongoing exposure pathway of PFOA to the background population.

Introduction

Perfluorocarboxylates (PFCAs) have been manufactured since 1947 initially by electrochemical fluorination (ECF) and, since the 1970s, also by telomerization (1). Manufacture of PFCAs using the ECF process was phased out between 2000 and 2002, but manufacture by telomerization is ongoing. The main use of PFCAs is as processing aids in fluoropolymer manufacture, but historically they have had a wide range of other minor uses (1). Direct industrial emissions are estimated to be the major source of PFCAs to the environment (1), although PFCAs are also breakdown products of “precursor” compounds such as fluorotelomer alcohols (FTOHs) and perfluoroalkyl sulfonamido alcohols (FOSE-OHs) (2–10) and processing impurities in commercial products (1). PFCAs are of scientific and regulatory interest as they are a class of

extremely persistent chemicals (11) that have been found globally in wildlife and in humans (12). Toxicology studies of PFOA in rodents have shown increased incidence of liver, pancreas, and testicular tumors in rodents although epidemiological studies indicate a lack of carcinogenic activity in humans (13). An observed negative association between PFOA serum concentration and birth weight has recently generated interest for the developmental toxicology of PFOA (14). However, further research is needed to determine if noncausal physiological associations could explain the alterations in fetal growth indicators (15).

Although the presence of PFCAs in human serum has generated a great deal of attention and environmental research, many questions remain about the pathways of human contamination (16). Source elucidation of PFCAs in humans has proven to be complicated because concentrations of PFCAs present in humans represent the integration of PFCA exposure from a vast number of pathways over an extended period of time. Here we critically evaluate the current published literature on human exposure to PFCAs with the aim of providing new insights into the pathways of human exposure. The review is divided into three sections which focus on (i) concentrations and trends in human sera, (ii) exposure estimates from concentrations of PFCAs in exposure media and, (iii) pharmacokinetics and pharmacokinetic modeling. Due to the prevalence of data, emphasis is placed on PFOA.

Concentrations, Homologue Patterns and Temporal Trends in Human Sera

PFCAs are distributed in extracellular space and are slowly eliminated in humans (17–20) with an estimated elimination half-life of 3.5 years (20). PFOA and occasionally longer chain homologues (C9 – C11) have been measured in human sera (17, 21–49), plasma (50–54), and whole blood (55–63). Concentrations of PFOA in serum or plasma and whole blood display an approximate 2:1 ratio (17) making measurements in these matrices comparable after correction. Some studies have also reported the presence of PFOA in liver, muscle, and adipose tissue (24, 58).

Analyzed concentrations of PFOA in sera from various populations can be divided into three categories based on the type of exposure (Figure 1). The highest serum concentrations of PFOA (median concentrations 1100–1300 $\mu\text{g/L}$) have been measured in ammonium perfluorooctanoate (APFO) production workers (22, 26). Elevated serum concentrations of PFOA (mean 27.4–423 $\mu\text{g/L}$) in nonoccupationally exposed populations have been reported from Little Hocking, U.S. (48) and Arnsberg, Germany (52) where the

* Corresponding author e-mail: ian.cousins@itm.su.se.




FIGURE 1. Human serum concentrations of PFOA ($\mu\text{g/L}$) in populations of (a) occupational exposure, (b) point source exposure, and (c) background exposure. Solid bars depict arithmetic mean concentrations and the min-max range is given by the error bars. Whole blood concentrations have been adjusted to serum concentration by multiplying with a factor of 2. References of the individual studies are given in square brackets on the x-axis.

use of contaminated soil conditioner and a fluoropolymer production site, respectively, were identified as sources of contamination. Systematic analysis showed that drinking water was the primary predictor of PFOA serum concentrations in Arnsberg and Little Hocking. However, in the majority of serum samples from the general population in different parts of the world, the concentrations of PFOA are reasonably consistent with only small interregional differences (Figure 1). For example, cross-continental biomonitoring studies of children, teenagers, adults, and elderly populations in the U.S. showed that PFOA concentrations in sera are log normally distributed with geometric mean values of 4–5 $\mu\text{g/L}$ (24, 25, 28, 34), with most other studies from industrialized countries in the rest of the world reporting mean concentrations between 2 and 8 $\mu\text{g/L}$ (Figure 1). Notable exceptions from this trend include high median concentrations of PFOA in populations from Korea (male 36.8; female 30.9 $\mu\text{g/L}$), Poland (male 20.5, female 23.2 $\mu\text{g/L}$), New York (25.2 $\mu\text{g/L}$), and Kentucky (male 38.1, female 20 $\mu\text{g/L}$) (40) as well as low concentrations in Peruvian citizens (75% of the samples <0.1 $\mu\text{g/L}$) (30). The comparable concentrations of PFOA in Brazil (10 $\mu\text{g/L}$), Colombia (6.2 $\mu\text{g/L}$) (40), and Australia (7.6 $\mu\text{g/L}$) (44) with those of Europe and North America are interesting because the Southern Hemisphere has no known manufacturing sources of PFOA (1) and background concentrations in the environment are much lower (45). Although, there are concerns about the accuracy and reproducibility in analytical methods for PFCAs, whole blood and serum measurements have been shown to be relatively reliable in an interlaboratory comparison (65).

The homologue pattern in human sera is typically PFOA > PFNA > PFDA (perfluorodecanoate), with PFCAs of 10 carbons or more being detected infrequently (Figure 2) (21, 23, 28–33, 44, 56, 59–62). A qualitative comparison of the homologue pattern in blood and other tissues of wildlife (12, 64–69) reveals that the dominance of PFOA is with a few exceptions (e.g., in sea turtles (72), sea otters (73), eel and

flounder (74) distinctive to humans (12). The homologue pattern in terrestrial and marine mammals, various fish, and bird species displays a characteristic dominance of odd-carbon chain lengths (e.g., C9, C11), which are at higher concentrations than the next shortest even-carbon homologues (e.g., C8, C10) (12, 64–69). Although wildlife samples comprise liver, egg, and whole body homogenates in addition to whole blood and serum samples, the PFCA profile is fairly consistent regardless of sampled tissue (70). The PFNA > PFOA pattern in biota has been attributed to increasing bioconcentration and biomagnification with PFCA chain length (70, 71). The dominance of PFOA in human serum leads to two different hypotheses: (i) the pathways of human exposure to PFOA are different from those of wildlife, i.e., not related to food chain bioaccumulation/biomagnification, and/or (ii) humans display different pharmacokinetics (e.g., different relative retention pattern of PFCAs with increasing chain lengths in humans relative to wildlife).

Although it is common to observe a positive correlation between serum concentrations and age for persistent compounds with relatively long elimination half-lives, no clear trend with age has been observed for PFOA (24, 25, 28). The lack of a trend may be due to (i) efficient *in utero* transfer of PFOA (27, 54, 55, 76), (ii) a higher body-weight-normalized exposure to infants, toddlers and children (78), (iii) increasing exposure levels over time in proportion to increasing production (35), or a combination of these factors. In contrast to age, a significant difference in PFOA serum concentrations has been determined with respect to sex and ethnicity (31, 53, 54, 56, 79), which implies that genetic variability or life-style factors affect either the exposure or pharmacokinetics of PFOA.

A number of studies analyzing PFCAs in human sera have reported a significant correlation between PFOS and PFOA in North America and Europe ($0.05 > p\text{-values} > 0.0001$, Table S2) (27–29, 32, 34–37, 40, 42, 53, 54, 59). The observed correlation between PFOS and PFOA is interesting because these com-




FIGURE 2. Human serum concentrations ($\mu\text{g/L}$) of PFOA, PFNA, PFDA, and PFUnA in background exposed populations. Solid bars represent arithmetic mean concentrations and the min–max range is given by the error bars. Whole blood concentrations have been adjusted to serum concentration by multiplying with a factor of 2. References of the individual studies are given in square brackets on the x-axis.

pounds cannot convert directly to each other, nor do they have a major source to the environment in common according to global emission inventories (1, 80). PFOS and other products based on POSF-based chemistry have been produced exclusively by the ECF process and furthermore PFOS is not a breakdown product of fluorochemicals produced by any other process. Releases of PFOA, however, result from both the ECF and telomerization production processes. PFOA is an intended end product and a residual impurity in products from both the main production processes, and both telomer- and POSF-based precursors that degrade to form PFOA in the environment. Pathways that could give rise to a human co-exposure of PFOA and PFOS in North America, Europe, and Australia include (i) microbial or atmospheric degradation (8, 9) of POSF-based precursors and subsequent coexposure to both PFOA/PFOS via environmental pathways, (ii) exposure to POSF-based precursor compounds that are metabolized to both PFOA and PFOS, or (iii) exposure to residual amounts of PFOA and PFOS from POSF-based consumer products (1). Regarding the second suggested pathway it should be noted that biotransformation studies have not identified PFCA metabolites from *N*-ethyl-sulfonamidoethanol (*N*-EtFOSE) in rats (5, 6) and *N*-ethyl-perfluorooctanesulfonamide (*N*-EtPFOSA) in trout (7). However, degradation products of POSF-based polyfluoroalkyl phosphate surfactants (PAPS) and polymeric materials remain to be identified.

De Silva and Mabury (21) proposed quantitative analysis of linear and branched carbon chain isomers in serum as a direct tool for source elucidation. ECF-produced PFCAs have been found to contain 20–30% of branched chain isomers, whereas telomerization produces >99% linear chain PFCAs (21). Thus, an observed dominance of linear PFOA in Canadian human sera was suggested to be evidence for a dominance of fluorotelomer-based sources. However, studies showing a preferential elimination of branched compared

to linear PFOA isomers in human (17), and animal studies (18) indicate that branched:linear ratios of PFOA isomers need to be corrected for the differences in pharmacokinetic behavior before being used for exposure interpretation.

Temporal trends of serum in the U.S. population have shown an increase of PFOA concentrations between 1974 and 1989 (from 2.2 to 5.5 $\mu\text{g/L}$) after which concentrations leveled off between 1989 and 2001 (5.5–4.2 $\mu\text{g/L}$) (27). A decrease in serum concentrations of PFOA has been observed in the U.S. and Norway starting at around year 2000 with estimated disappearance half-lives of approximately 4.4 years (29, 35–37, 76, 77) (Figure 3a), which are slightly longer than the elimination half-life reported in retired workers (20). Given the continued production of APFO (Figure 3b), the environmental persistence of PFOA and absence of declining concentration trends in wildlife (67–69), the rapid response in human serum likely reflects a decrease in consumer product exposure. As ECF was the sole manufacturing process used for PFOS, the simultaneous decrease of PFOA and PFOS (Figure S1) (29, 35–37, 76) indicates that exposure pathways of these compounds were historically linked. Only POSF production has reduced since 2000 (81) (Figure 3b) (fluorotelomer and APFO production has remained constant) indicating a link between the 2000–2002 POSF phase-out and the decline of PFOS/PFOA in human serum. In comparison to PFOA and PFOS, it is interesting to note that PFNA and PFDA serum concentrations have been constant or slightly increasing (29, 76, 77) after year 2000 (Figures S2 and S3), indicating an ongoing exposure for these compounds and/or that serum half-lives are much longer. Contrary to the clear downward trends in the Northern Hemisphere, preliminary data from Australian serum samples indicate no decrease in PFOA concentrations between 2002/2003 (44) and 2006/2007 (82).




FIGURE 3. (a) Temporal trends of PFOA in human serum ($\mu\text{g/L}$) from pooled serum, Norway (77), NHANES, U.S. (29), and Red cross blood donors (35, 37). The temporal trend in infant blood spots ($\mu\text{g/L}$) (76) is plotted on the second y-axis. This was done to facilitate comparison of time trends. The blood spot concentrations were about a factor of 5 lower than serum concentrations presumably because (i) they were blood and not serum samples (approximately a factor of 2 lower), and (ii) they were from infants rather than adults. (b) Estimated production of POSF, APFO and Fluorotelomer A, (tonnes/annum) (1, 87).

Taken as a whole, the inter-regionally consistent serum concentrations of PFOA, characteristic homologue pattern (PFOA > PFNA), and rapid response to production changes indicate that exposure pathways of PFOA to humans are different from those to wildlife. Furthermore, the PFOS:PFOA correlation and similar time trend for these compounds indicate that human serum levels are influenced by a historical POSF-based exposure pathway.

Exposure Estimates, Homologue Patterns and Temporal Trends in Exposure Media

Measured concentrations in exposure media combined with estimated exposure factors provide a comparison of the relative importance of different exposure pathways. To date, measurements of multiple exposure media for a single population are not available. However, a compilation of estimated daily intakes (83, 85–91) for populations of typical background exposure, elevated water concentrations, and occupational exposure

illustrate the relative importance of different exposure pathways (Figure 4). In line with previous assessments (78), dietary intake as determined by a duplicate serving study in Germany (87) is estimated to be the dominant pathway of exposure for the background exposed population (Figure 4a). House dust (86) and “background” drinking water (88) are estimated to be minor exposure pathways. Drinking water concentrations of PFOA display large interregional variability (48, 89, 91, 93–95) from below 0.32 (88) up to 7200 ng/L (48). The relative importance of drinking water as an exposure medium becomes significant at concentrations of 40 ng/L (90), and dominates exposure at 519 ng/L (91) (Figure 3b and c). Although limited data are available for occupational exposure, ambient air concentrations of PFOA ($1 \mu\text{g}/\text{m}^3$) in the vicinity of a fluoropolymer manufacture facility (92) suggest that inhalation is a major route of exposure for fluorochemical workers, although dermal contact and incidental ingestion (not quantified here) may also contribute (26) (Figure 4d).




FIGURE 4. Pie charts displaying a compilation of the estimated daily intakes of male adults (D) and relative importance of exposure pathways from separate studies. Each pie chart represents an exposure scenario representative of (a) background concentrations in drinking water (1.3 ng/L); (b) elevated concentrations in drinking water (40 ng/L); (c) point sources of drinking water contamination (519 ng/L); (d) occupationally exposed individuals (indoor air concentrations $1 \mu\text{g}/\text{m}^3$). References of the individual studies are given in square brackets in the legends of each chart.

Previous studies have identified PFOA as a metabolite of precursor compounds such as fluorotelomer alcohols (FTOHs) and telomer-based (PAPS) (21, 84). Exposure to FTOHs has been estimated to occur primarily via food packaging materials (96, 97), indoor air (85, 98), and house dust (86). As displayed in Figure 4, the biotransformation of FTOHs was estimated to be of minor significance for the daily intakes of PFOA (83) due to the small fraction of 8:2 FTOH forming PFOA in laboratory *in vivo* and *in vitro* experiments (2–4). A recent study showing substantial amounts of PAPS in pooled human serum from the U.S. highlights the need for inclusion of these compounds in exposure assessments (99). An improved pharmacokinetic understanding of FTOHs and PAPS metabolism and measured concentrations in exposure media (e.g., food) would help to evaluate the relative importance of precursor compounds.

Estimated exposures extrapolated from exposure media (78, 83) in Figure 4 may be limited by the variability in environmental concentrations, consumer behavior, product formulations, or as discussed later, unknown exposure pathways. Trudel et al. (78) illustrated that uncertainty in measured concentrations of food items had the highest influence on variability in estimated intakes. Due to the challenges of overcoming matrix effects in the analysis of foodstuffs, most current reported concentrations of PFOA in food are below the current limit of quantification of 0.1–1 ng/g. Infrequent detections of PFOA have been reported in a variety of food items including fish (100–103), meat (100), potatoes (105), fast food (100), milk, apples, green beans, bread, and eggs (see Table S8). Clearly, improved analytical data are needed for food items as well as an improved understanding of the sources of PFCA contamination in food.

The relationships between PFCA homologues in food items and human serum will provide valuable insights into the sources of food contamination, but unfortunately there is currently a paucity of measurements of PFCA profiles in food. It might be expected that if meat and fish contaminated via bioaccumulation in food webs were the major source of

PFCA in food then a PFNA \geq PFOA homologue pattern would be expected in total diet intake (100–102). Interestingly, a Japanese duplicate diet study (104) measured PFOA at concentrations similar to those of Fromme et al. (87), whereas PFNA was not detected in any of the samples. The different homologue pattern of individual wild food items (100–102) compared to duplicate food samples (104) provides a preliminary indication that bioaccumulative pathways related to environmental contamination are not the major source of PFOA in food. Studies investigating the importance of food chain pathways found positive correlations between the consumption of seafood (61) and whale meat (47) and serum concentrations of PFNA, PFDA, PFUnA, and perfluorododecanoate (PFDoA). However, for PFOA only weak correlations were observed between serum concentrations and consumption of fish (61) and red meat (75) (other homologues were not reported for red meat). These studies indicate that exposure pathways to longer chain homologues ($>C8$) are distinct from those for PFOA. Potential pathways that could result in a dominance of PFOA in the total diet include the uptake in high water content crops (88, 91, 93–95, 106) or migration from food packaging paper (96, 97). A recent study by Stahl et al. (106) demonstrated that soil-to-plant uptake of PFOA occurs in maize, oats, wheat, and potatoes with concentrations in the ears, grain, and tuber reaching 0.1–3% of the concentration in soil. Recently, it has been shown that agricultural soils have been contaminated with PFOA via application of sewage sludge, providing a transfer mechanism of commercially produced food that may be different from that of wild food (107).

A significant drawback in understanding exposure is the lack of comprehensive time trends for PFCA concentration data in exposure media. There are, nevertheless, some indications of a decline in consumer product-related exposure pathways. House dust samples collected in 2000–2003 from American (86) and Canadian (108) homes contain higher PFOA concentrations compared to more recently collected samples from Japan (109), Sweden (110), and Germany (111).

Although the difference among the studies could be due to regional and methodological differences, higher concentrations of PFOA in archived carpet protectors (112) compared with currently used alternatives (89) and/or a larger historical market share of POSF-based products could explain the higher concentrations in archived samples. A positive correlation ($r > 0.82$, $p < 0.0001$) between PFOA and PFOS in house dust (86) indicates that POSF-based products may be a common exposure pathway of these chemicals to the indoor environment. A sharp decrease in concentrations of ($8.3 - <0.1$ ng/g) of *N*-ethyl perfluorooctanesulfonamide (*N*-EtPFOA) in Canadian fast food composites has been reported between 1994 and 2002 (113), as a hypothesized consequence of changes in food paper formulations. Although *N*-EtPFOA is not believed to metabolize to PFOA (5–8), POSF-based food packaging materials contained a suite of residual perfluoroalkyl compounds that were not measured in the study (113). Interestingly, the initiation year of the decrease (1994) indicates that certain consumer-based pathways of PFCAs disappeared before completion of the ECF phase-out in 2000–2002. To address the paucity of temporally resolved exposure data, analysis of archived exposure media (e.g., food, house dust, and consumer products) for PFCAs and related compounds is needed.

Pharmacokinetics and Pharmacokinetic Modeling of PFCAs

The pharmacokinetic behavior of PFOA, and to a lesser extent other PFCAs, have been studied in laboratory animals and humans. Oral bioavailability of APFO has been determined to be between 66 and ~100% (20, 114). Efficient uptake via inhalation of APFO which is rapidly converted to PFOA *in vivo* has been observed, although no quantitative uptake efficiencies have been reported (115, 116). Dermal exposure studies have shown that PFOA can penetrate the skin, albeit with a low absorption efficiency (117). The elimination half-lives of PFOA reported in rats (114, 118–122), dogs (118), mice (123), and cynomolgus monkeys (124) display a large interspecies variation (<1 day for female rats to approximately 1 month for monkeys) and gender specificity. In comparison, estimated serum half-lives for PFOA in human serum are much longer with a geometric mean value of 3.5 years (range 1.5–9.1 years) reported for retired workers (20). The half-life of serum elimination was not associated with initial concentrations, gender, age, or time elapsed between the retirement and first blood collection (20). Scaling of animal derived half-lives (114, 118–122) with hepatic blood flow rates (125, 126) cannot replicate empirically determined elimination half-lives of PFOA in humans (20). Recently, renal resorption mediated by organic anion transporters (OATs) was proposed as a mechanism that could explain the observed species and gender based differences in elimination half-lives (127). As previously mentioned, the relative retention of PFCAs with varying chain lengths has an important implication for the interpretation of the PFCA homologue patterns in human serum (PFOA > PFNA). In pharmacokinetic studies, renal clearance rates have generally been observed in the order PFOA > PFNA > PFDA (121, 123), which is also consistent with the current understanding of elimination behavior in wildlife (71). However, measured tissue distributions and elimination half-lives of PFCAs other than PFOA are still missing in humans.

A common approach to characterizing the chemical concentration in a specific physiological compartment such as blood involves the use of physiologically based pharmacokinetic (PBPK) modeling with a detailed description of the major groups of tissues or organs together with the simulation of the arterial and venous circulatory system (128). Under steady-state conditions when intakes are fairly constant with

time, the rate of chemical intake matches the rate of chemical elimination and the concentrations in the various tissues or physiological compartments do not change significantly with time, creating constant concentration ratios among the various sites of chemical deposition in the body. In these cases, relatively simple models can be developed based on noncompartmental, one-compartment, or two-compartment concepts. A one-compartment steady-state pharmacokinetic model (eq 1) has recently been applied to relate internal concentrations of PFOA in humans to estimated daily intakes (129). It is thought to be particularly applicable for persistent compounds such as PFOA and for the low concentrations typical of environmental exposure.

$$C_{ss} = \frac{T_{1/2}D}{0.693 \cdot V_D} \quad (1)$$

Where the steady-state serum concentration (C_{ss}) is a function of the total elimination half-life ($T_{1/2}$) (discussed above), body weight normalized daily internal intake or dose (D), and fractional volume of distribution (V_D) (discussed below). By rearrangement, the model can also be used to back-calculate D if C_{ss} is available (aka “reverse dosimetry”).

Estimates of V_D have been obtained from single-dose studies as well as multiple dosing studies; primarily in rodents and primates (see Table S3). The estimated values depend on experimental design (type of dosing) and the model used for analysis. Single-dose studies have reported values of V_D of 210–340 mL/kg (114, 121) in rats and 181–198 mL/kg in monkeys (124), indicating that single doses of PFOA are distributed primarily in the blood (121). Based on repeated dose studies in monkeys, V_D values of 1260–6340 mL/kg have been estimated for subchronic dosing (124). Thus, V_D values calculated from steady-state dosing of PFOA are approximately an order of magnitude higher than those derived from a single dose. The increase in V_D following continuous dosing indicates that PFOA distributes to extracellular fluids and highly perfused organs such as liver and kidney in addition to blood.

In the following model evaluation a V_D derived from subchronic dose studies (i.e., 3.6 L/kg) was used as it represents long-term exposure of humans more realistically. As previously shown in this review, drinking water concentrations exceeding 500 ng/L constitute the major exposure pathway for nonoccupationally exposed populations (Figure 3c). Hence, measured serum concentrations from subpopulations exposed to elevated drinking water concentrations can be used to evaluate model accuracy. To represent the variability and uncertainty in V_D , $T_{1/2}$, and C_{ss} , a probabilistic approach was used to simulate the expected daily intakes associated with measured serum concentrations (parameterization in SI). Figure 5a shows the back-calculated internal daily intake from serum concentrations compared to estimated internal daily intakes derived from drinking water concentrations. The pharmacokinetic model estimates the intakes within a factor 1.5–5.5 for populations in the Arnsberg (52, 91), Little Hocking (48, 49), and Lubeck (49, 51) regions. Noteworthy is the large uncertainty in predicted daily intakes for Little Hocking (2005–2006) and Lubeck (2005–2006). A sensitivity analysis showed that serum concentrations had the highest influence on back-calculated daily intakes from Little Hocking and Lubeck (Figures S7–S8), indicating that duration and frequency of drinking water exposure displayed large variability. The model overprediction in Arnsberg and Little Hocking (2002–2005) suggests that the value of V_D is too high (Figures S5 and S6).

We can also use the same model parameterization to back-calculate internal daily intakes of PFOA from serum concentrations for a background exposed population in Bavaria, Germany in 2005 (53) for which daily dietary intakes of PFOA




FIGURE 5. Comparison of estimated daily intakes (D) of PFOA using measurements of exposure media and the one-compartment pharmacokinetic relationship for (a) exposure to point source contaminated drinking water and (b) background exposure to food. Solid bars represent median values and the error bars represent the 10th and 90th percentile, respectively. References of the individual studies used for making the estimates of daily intake are given in square brackets in the chart legends. Calculations are explained in more detail in the Supporting Information.

are also available (87). The measured median daily internal intake of PFOA from food (87) was determined to be a factor of 2.6 lower than back-calculated internal daily intakes from matched serum measurements (Figure 5b) (53). It should be noted that similar calculations have recently been made. Trudel et al. (78) concluded that estimated internal daily intake from multiple pathways (3 ng/(kg day) median) were within the range of back-calculated daily internal intakes from serum concentrations for the background population. However, Fromme et al. (87) concluded that the daily internal intake of PFOA (derived only from dietary intake) was a factor of 6 higher than the back-calculated internal daily intake from serum concentrations despite the similar estimated intake (2.9 ng/(kg day)). The discrepancy of these two earlier studies is primarily explained by the chosen value of V_D (0.2 L/kg in ref 87 and median of 3.6 L/kg in ref 78) illustrating the importance of correct parametrization of eq 1.

Despite the incomplete understanding of the pharmacokinetic behavior of PFCAs, the one-compartment model, if correctly parameterized, provides an estimation within a factor of 1.5–5.5 of the internal daily intakes. The disagreement between the two estimates could be a result of

uncertainty in model input parameters, primarily V_D . As serum concentrations of PFOA from 2005 reflect both ongoing and historical pathways the overprediction of daily intakes from the pharmacokinetic model may be due to the inappropriateness of the steady-state assumption. The model evaluations (Figure 5) provide further supporting evidence that dietary exposure is a major ongoing exposure pathway of PFCAs, except near contaminated drinking water sources, where drinking water exposure dominates.

Concluding Remarks

Research presented in this review taken as a whole supports the hypothesis that contemporary serum concentrations of PFOA in humans are influenced by an unknown historical exposure source derived from production of POSF-based consumer products prior to the 2000–2002 phase-out by the major manufacturer. The estimates made here and elsewhere that food dominates the current estimated daily intakes of PFOA are primarily supported by two separate duplicate diet studies performed in Germany (87) and Japan (104), respectively. Nevertheless, improved analytical methods and

larger analyzed sample sets within the European research collaboration, PERFOOD (130), will soon provide more accurate estimates of the daily intake via food as well as investigate how the food is being contaminated (i.e., the relative importance of environmental contamination versus contamination from food processing/packaging).

As the fluorine industry has announced a commitment to the phase-out of eight carbon perfluoroalkyl chemistry by 2015 (131), future human exposure to PFOA will depend progressively more on legacy sources of contamination in the environment and less on consumer product-derived exposure pathways. Therefore the currently observed rate of decline of PFOA in human serum since the phase-out of POSF-based products by the major manufacturer will probably not be sustained much longer as environmentally derived exposure pathways become relatively more important. There will, however, probably still be a continued albeit slower decline as the concentrations of environmentally contaminated food will also decline in proportion to the estimated future declines in environmental concentrations in temperate/source regions (132) where much of the global food is produced. It has been hypothesized (132) that (i) the worlds oceans will contain the majority of the legacy releases of PFOA, (ii) once releases cease the levels in the surface oceans will gradually become approximately the same as ocean water becomes well-mixed with respect to PFOA, and (iii) PFOA will only slowly be removed from the surface oceans by vertical transport to the deep oceans. Therefore, into the far future humans will continue to receive a fairly constant, low-level exposure to PFOA from the consumption of marine fish and other seafood.

Acknowledgments

We thank E.I. DuPont de Nemours & Co., Inc. for an unrestricted grant which helped fund the Ph.D. studies of R.V. We thank Mark H. Russell, Robert C. Buck, Martin Scheringer, David Trudel, Urs Berger, Merle Plassmann, and James Armitage for providing valuable comments.

Supporting Information Available

Summary of PFOS:PFOA correlations in human serum; temporal trends of PFOS, PFNA, and PFDA in human serum; tables of concentrations of PFCAs and precursors in exposure media; an explanation of the derivation of the one-compartment pharmacokinetic model; and further details of the uncertainty analysis including contribution to variance plots. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Literature Cited

- Prevedouros, K.; Cousins, I. T.; Buck, R. C.; Korzeniowski, S. H. Sources, Fate and Transport of Perfluorocarboxylates. *Environ. Sci. Technol.* **2006**, *40*, 32–44.
- Martin, J. W.; Mabury, S. A.; O'Brien, P. Metabolic products and pathways of fluorotelomer alcohols in isolated rat hepatocytes. *Chem.-Biol. Interact.* **2005**, *155*, 165–180.
- Fasano, W. J.; Carpenter, S. C.; Gannon, S. A.; Snow, T. A.; Stadler, J. C.; Kennedy, G. L.; Buck, R. C.; Korzeniowski, S. H.; Hinderliter, P. M.; Kemper, R. A. Absorption, distribution, metabolism and elimination of 8:2 fluorotelomer alcohol in the rat. *Toxicol. Sci.* **2006**, *91*, 341–355.
- Nabb, D. L.; Szostek, B.; Himmelstein, M. W.; Mawn, M. P.; Gargas, M. L.; Sweeny, L. M.; Stadler, J. C.; Buck, R. C.; Fasano, W. J. In-vitro metabolism of 8–2 fluorotelomer alcohol: Interspecies comparisons and metabolic pathway refinement. *Toxicol. Sci.* **2007**, *100*, 333–344.
- Tomy, G. T.; Tittlemier, S. A.; Palace, V. P.; Budakowski, W. R.; Braekvelt, E.; Brinkworth, L.; Friesen, K. Biotransformation of N-Ethyl perfluorooctanesulfonamide by rainbow trout (*Oncorhynchus mykiss*) liver micromes. *Environ. Sci. Technol.* **2004**, *38*, 758–762.
- Xu, L.; Krenitsky, D. M.; Seacat, A. M.; Butenhoff, J. L.; Anders, M. W. Biotransformation of N-Ethyl-N-(2-hydroxyethyl)perfluorooctanesulfonamide by rat liver microsomes, cytosol, and slices and by expressed rat and human cytochromes P450. *Chem. Res. Toxicol.* **2004**, *17*, 767–775.
- Xu, L.; Krenitsky, D. M.; Seacat, A. M.; Butenhoff, J. L.; Tephly, T. R.; Anders, M. W. N-Glucuronidation of perfluorooctanesulfonamide by human, rat, dog and monkey liver microsomes and by expressed rat and human UDP Glucuronosyltransferases. *Drug Metab. Dispos.* **2006**, *34*, 1406–1410.
- D'Eon, J. C.; Hurley, M. D.; Wallington, T. J.; Mabury, S. A. Atmospheric Chemistry of N-methyl perfluorobutane sulfonamidoethanol, C₄F₉SO₂N(CH₃)CH₂CH₂OH: Kinetics and mechanism of reaction with OH. *Environ. Sci. Technol.* **2006**, *40*, 1862–1888.
- Martin, J. W.; Ellis, D. A.; Mabury, S. A.; Hurley, M. D.; Wallington, T. J. Atmospheric chemistry of perfluoroalkanesulfonamides: Kinetic and product studies of the OH radical and Cl atom initiated oxidation of N-ethyl perfluorobutanesulfonamide. *Environ. Sci. Technol.* **2006**, *40*, 864–872.
- Lange, C. C. *The aerobic biodegradation of N-EtFOSE alcohol by the microbial activity present in municipal wastewater treatment sludge: Biodegradation study report*; 3MProject ID: LIMS E00-2252; 3M Company: St. Paul, MN, 2000.
- Kissa, E. *Fluorinated Surfactants and Repellents*, 2nd ed.; Marcel Dekker: New York, 2001.
- Houde, M.; Martin, J. W.; Letcher, R. J.; Solomon, K. R.; Muir, D. C. G. Biological monitoring of perfluoroalkyl substances: A review. *Environ. Sci. Technol.* **2006**, *40*, 3463–3473.
- Kennedy, G. L.; Butenhoff, J. L.; Olsen, G. W.; O'Connor, J. C.; Seacat, A. M.; Perkins, R. G.; Biegel, L. B.; Murphy, S. R. The toxicology of perfluorooctanoate. *Crit. Rev. Toxicol.* **2004**, 351–384.
- Apelberg, B. J.; Witter, F. R.; Herbstman, J. B.; Calafat, A. M.; Halden, R. U.; Needham, L. L.; Goldman, L. R. Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in Relation to Weight and Size at Birth. *Environ. Health Perspect.* **2007**, *115*, 1670–1676.
- Andersen, M. E.; Butenhoff, J. L.; Chang, S. C.; Farrar, D. G.; Kennedy, G. L.; Lau, C.; Olsen, G. E.; Seed, J.; Wallace, K. B. Perfluoroalkyl acids and related chemistries-Toxicokinetics and modes of action. *Toxicol. Sci.* **2008**, *102*, 3–14.
- Renner, R. PFOA in people. *Environ. Sci. Technol.* **2007**, *41*, 4497–4500.
- Ehresman, D. J.; Froelich, J. W.; Olsen, G. W.; Chan, S. C.; Butenhoff, J. L. Comparison of whole blood, plasma and serum matrices for the determination of perfluorooctanesulfonate (PFOS), perfluorooctanoate (PFOA) and other fluorochemicals. *Environ. Res.* **2007**, *103*, 176–184.
- Loveless, S. E.; Finlay, C.; Everds, N. E.; Frame, S. R.; Gillies, P. J.; O'Connor, J. C.; Powley, C. R.; Kennedy, G. L. Comparative responses of rats and mice exposed to linear/branched, linear, or branched ammonium perfluorooctanoate (APFO). *Toxicology* **2006**, *220*, 203–217.
- Hundley, S. G.; Sarraf, A. M.; Kennedy, G. L. Absorption, distribution and excretion of Ammonium Perfluorooctanoate (APFO) after oral administration to various species. *Drug Chem. Toxicol.* **2006**, *29*, 137–145.
- Olsen, G. W.; Burris, J. M.; Ehresman, D. J.; Froelich, J. W.; Seacat, A. M.; Butenhoff, J. L.; Zobel, L. R. Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate and perfluorooctanoate in retired fluorochemical workers. *Environ. Health Perspect.* **2007**, *115*, 298–305.
- De Silva, A. O.; Mabury, S. A. Isomer distribution of perfluorocarboxylates in human blood: Potential correlation to source. *Environ. Sci. Technol.* **2006**, *40*, 2903–2909.
- Olsen, G. W.; Burris, J. M.; Burlew, M. M.; Mandel, J. H. Plasma cholestokinin and hepatic enzymes, cholesterol and lipoproteins in ammonium perfluorooctanoate production workers. *Drug Chem. Toxicol.* **2000**, *23*, 603–620.
- Miyake, Y.; Yamashita, N.; So, K. M.; Rostkowski, P.; Taniyasu, S.; Lam, P. K. S.; Kannan, K. Trace analysis of total organic fluorine in human blood using combustion ion chromatography for fluorine: A mass balance approach for the determination of known and unknown organofluorine compounds. *J. Chromatogr.* **2007**, *115*, 214–221.
- Olsen, G. W.; Hansen, K. J.; Stevensson, L. A.; burris, J. M.; Mandel, J. H. Human donor liver and serum concentrations of perfluorooctane sulfonate and other perfluorochemicals. *Environ. Sci. Technol.* **2003**, *37*, 888–891.
- Olsen, G. W.; Church, T. R.; Miller, J. P.; Burris, J. M.; Hansen, K. J.; Lundberg, J. K.; Armitage, J. B.; Herron, R. M.; Medhizadehkashi, Z.; Nobilet, J. B.; O'Neill, E. M.; Mandel, J. H.;

- Zobel, L. R. Perfluorooctanesulfonate and Other Fluorochemicals in the Serum of American Red Cross Adult Blood Donors. *Environ. Health Perspect.* **2003**, *111*, 1892–1901.
- (26) Olsen, G. W.; Logan, P. W.; Hansen, K. J.; Simpson, C. A.; Burris, J. M.; Burlew, M. M.; Vorarath, P. P.; Venkateswarlu, P.; Schumpert, J. C.; Mandell, J. H. An Occupational Exposure Assessment of a Perfluorooctanesulfonyl Fluoride Production Site: Biomonitoring. *AIHA J.* **2003**, *64*, 651–659.
- (27) Apelberg, B. J.; Goldman, L. R.; Calafat, A. M.; Herbstman, J. B.; Kuklenyik, Z.; Heidler, J.; Needham, L. L.; Halden, R. U.; Witter, F. Determinants of fetal exposure of perfluoroalkyl substances in Baltimore, Maryland. *Environ. Sci. Technol.* **2007**, *41*, 3891–3897.
- (28) Calafat, A. M.; Kuklenyik, Z.; Reidy, J. A.; Caudill, S. P.; Needham, L. L. Serum concentrations of polyfluoroalkyl compounds in the U. S. population: Data from the National Health and Nutrition Examination Study (NHANES) 1999–2000. *Environ. Sci. Technol.* **2007**, *41*, 2237–2242.
- (29) Calafat, A. M.; Wong, L.-Y.; Kuklenyik, Z.; Reidy, J. A.; Needham, L. Polyfluoroalkyl chemicals in the U.S. population: Data from the National Health and Nutrition survey (NHANES) 2003–2004 and comparisons to NHANES 1999–2000. *Environ. Health Perspect.* **2007**, *115*, 1596–1602.
- (30) Calafat, A. M.; Needham, L. L.; Kuklenyik, Z.; Reidy, J. A.; Tully, J. S.; Aguilar-Villalobos, M.; Naeher, L. P. Perfluorinated chemicals in selected residents of the American continent. *Chemosphere* **2005**, 490–496.
- (31) Calafat, A. M.; Kuklenyik, Z.; Caudill, S.; Reidy, J. P.; Needham, L. L. Perfluorochemicals in pooled serum samples from United States residents in 2001 and 2002. *Environ. Sci. Technol.* **2006**, *40*, 2128–2134.
- (32) Kuklenyik, Z.; Adam Reich, J.; Tully, J. S.; Calafat, A. M. Automated solid phase extraction and measurement of perfluorinated organic acids and amides in human serum and milk. *Environ. Sci. Technol.* **2004**, *38*, 3698–3704.
- (33) Kuklenyik, Z.; Needham, L. L.; Calafat, A. M. Measurement of 18 perfluorinated organic acids and amides in human serum using on-line solid-phase extraction. *Anal. Chem.* **2005**, *77*, 6085–6091.
- (34) Olsen, G. W.; Church, T. R.; Larson, E. B.; van Belle, G.; Lundberg, J. K.; Hansen, K. J.; Burris, J. M.; Mandel, J. H.; Zobel, L. R. Serum concentrations of perfluorooctanesulfonate and other perfluorochemicals in an elderly population from Seattle, Washington. *Chemosphere* **2004**, *54*, 1599–1611.
- (35) Olsen, G. W.; Huang, H. Y.; Helzlsouer, K. J.; Butenhoff, J. L.; Mandel, J. H. Historical comparison of perfluorooctanesulfonate, perfluorooctanoate and other fluorochemicals in human blood. *Environ. Health Perspect.* **2005**, *113*, 539–545.
- (36) Olsen, G. W.; Mair, D. C.; Reagen, W. K.; Ellefson, M. E.; Ehresman, D. J.; butenhoff, J. L.; Zobel, L. R. Preliminary evidence of a decline in perfluorooctanesulfonate (PFOS) and perfluorooctanoate (PFOA) concentrations in American Red Cross blood donors. *Chemosphere* **2007**, *68*, 105–111.
- (37) Olsen, G. W.; Mair, D. C.; Church, T. R.; Ellefson, M. E.; Reagen, W. K.; Boyd, T. M.; Herron, Z. M.; Noblietti, J. B.; Rios, J. A.; Butenhoff, J. L.; Zobel, L. R. Decline in perfluorooctanesulfonate and other perfluoroalkyl chemicals in American Red Cross adult blood donors, 2000–2006. *Environ. Sci. Technol.* **2008**, *42*, 4989–4995.
- (38) Olsen, G. W.; Church, T. R.; Hansen, K. J.; Burris, J. M.; Butenhoff, J. L.; Mandel, J. H.; Zobel, L. R. Quantitative Evaluation of Perfluorooctanesulfonate (PFOS) and Other Fluorochemicals in the Serum of Children. *J. Child. Health* **2004**, *2*, 53–76.
- (39) Ellefson, M. E.; Reagen, W. K. Definitive Study: Quantitative Screening of Commercial Lots of Human Serum Obtained in 2004 for Endogenous Fluorochemicals Using Solid Phase Extraction and Liquid Chromatography/Tandem Mass Spectrometry; 3M Project No. E05-0442, USEPA Administrative Record AR226-3579; Washington, DC, 2005.
- (40) Kannan, K.; Corsolini, S.; Falandysz, J.; Fillmann, G.; Senthil Kumar, K.; Lognathan, B.; Ali Mohd, M.; Olivero, J.; van Wouwe, N.; Ho Yang, J.; Aldous, K. M. Perfluorooctanesulfonate and related fluorochemicals in human blood from several countries. *Environ. Sci. Technol.* **2004**, *38*, 4489–4495.
- (41) Hansen, K. J.; Clemen, L. A.; Ellefson, M. E.; Johnson, H. O. Compound-specific, quantitative characterization of organic fluorochemicals in biological matrices. *Environ. Sci. Technol.* **2001**, *35*, 766–770.
- (42) Kubwabo, C.; Vais, N.; Benoit, F. M. A pilot study on the determination of perfluorooctanesulfonate and other perfluorinated compounds in blood of Canadians. *J. Environ. Monit.* **2004**, *6*, 540–545.
- (43) Harada, K.; Koizumi, A.; Norimitsu, S.; Inoue, K.; Yoshinaga, T.; Date, C.; Fujii, S.; Hachiya, N.; Hirose, I.; Koda, S.; Kusaka, Y.; Murata, K.; Omae, K.; Shimbo, S.; Takenaka, K.; Takeshita, T.; Todoriki, H.; Wada, Y.; Watanabe, T.; Ikeda, M. Historical and geographic trends of the increasing perfluorooctanoate and perfluorooctane sulfonate contamination in human serum in Japan. *Chemosphere* **2007**, *66*, 293–301.
- (44) Kärman, A.; Mueller, J. F.; van Bavel, B.; Harden, F.; Toms, L.-M. L.; Lindström, G. Levels of 12 perfluorinated chemicals in pooled Australian serum collected 2002–2003, in relation to age gender and region. *Environ. Sci. Technol.* **2006**, *40*, 3742–3748.
- (45) Yamashita, N.; Kannan, K.; Taniyasu, S.; Horii, Y.; Petrick, G.; Gamo, T. A global survey of perfluorinated acids in oceans. *Mar. Pollut. Bull.* **2005**, *51*, 658–668.
- (46) Kärman, A.; Ericson, I.; van Bavel, B.; Darnerud, P. O.; Aune, M.; Glynn, A.; Lignell, S.; Lindström, G. Exposure of perfluorinated chemicals through lactation: levels of matched human milk and serum and a temporal trend, 1996–2004, in Sweden. *Environ. Health Perspect.* **2007**, *115* (2), 226–230.
- (47) Weihe, P.; Kato, K.; Calafat, A. M.; Nielsen, F.; Wanigatunga, A. A.; Needham, L. L.; Grandjean, P. Serum concentrations of polyfluoroalkyl compounds in Faroese whale meat consumers. *Environ. Sci. Technol.* **2008**, *42*, 6291–6295.
- (48) Emmett, E. A.; Shofer, F. S.; Zhang, H.; Freeman, D.; Desai, C.; Shaw, L. M. Community exposure to perfluorooctanoate: Relationships between serum concentrations and exposure sources. *J. Occup. Environ. Med.* **2006**, *48*, 759–770.
- (49) West Virginia University School of Medicine. *C8 health project results*; December 2008; <http://www.hsc.wvu.edu/som/cmed/c8/results/C8AndC8SulfonateLevels/index.asp>.
- (50) Tao, L.; Kannan, K.; Aldous, K. M.; Mauer, M. P.; Eadon, G. A. Biomonitoring of perfluorochemicals in plasma of New York State personnel responding to the World Trade Center disaster. *Environ. Sci. Technol.* **2008**, 3472–3478.
- (51) Anderson-Mahoney, P.; Kotlerman, J.; Takhar, H.; Gray, D.; Dahlgren, J. Self-reported health effects among community residents exposed to perfluorooctanoate. *New. Solut.* **2008**, *18*, 129–143.
- (52) Hölzer, J.; Midash, O.; Rachfuss, K.; Kraft, M.; Reupert, R.; Angerer, J.; Kleeschulte, P.; Marschall, N.; Wilhelm, M. Biomonitoring of perfluorinated compounds in children and adults exposed to perfluorooctanoate (PFOA)-contaminated drinking water. *Environ. Health Perspect.* **2008**, *117*, 651–657.
- (53) Fromme, H.; Midash, O.; Twardella, D.; Angerer, J.; Boehmer, S.; Liebl, B. Occurrence of perfluorinated substances in an adult German population in southern Bavaria. *Int. Arch. Occup. Environ. Health* **2007**, *80*, 313–319.
- (54) Midash, O.; Schettgen, T.; Angerer, J. Pilot study on the perfluorooctanesulfonate and perfluorooctanoate exposure of the German general population. *Int. J. Hyg. Environ. Health* **2006**, *209*, 489–496.
- (55) Inoue, K.; Okada, F.; Ito, R.; Sasaki, S.; Nakajima, S.; Unno, A.; Saijo, Y.; Sata, F.; Yoshimura, Y.; Kishi, R.; Nakasawa, H. Perfluorooctane sulfonate (PFOS) and related perfluorinated compounds in human maternal and cord blood samples: Assessment of PFOS exposure in a susceptible population during pregnancy. *Environ. Health Perspect.* **2004**, *112*, 1204–1207.
- (56) Yeung, L. W. Y.; So, K. M.; Jiang, G.; Taniyasu, S.; Yamashita, N.; Song, M.; Wu, Y.; Li, J.; Giesy, J. P.; Guruge, K. S.; Lam, P. K. S. Perfluorooctanesulfonate and related fluorochemicals in human blood samples from China. *Environ. Sci. Technol.* **2006**, *40*, 715–720.
- (57) Guruge, K. S.; Taniyasu, S.; Yamashita, N.; Wijeratna, S.; Mohotti, K. M.; Seneviratne, H. R.; Kannan, K.; Yamanaka, N.; Miyazaki, S. Perfluorinated organic compounds in human blood serum and seminal plasma: a study of urban and rural tea workers in Sri Lanka. *J. Environ. Monit.* **2005**, *7*, 371–377.
- (58) Maestri, L.; Negri, S.; Ferrari, M.; Ghittori, S.; Fabris, F.; Danesino, P.; Imbriani, M. Determination of perfluorooctanoic acid and perfluorooctanesulfonate in human tissues by liquid chromatography/single quadrupole mass spectrometry. *Rapid Commun. Mass Spectrom.* **2006**, *20*, 2728–2734.
- (59) Ericson, I.; Gómez, M.; Nadal, M.; van Bavel, B.; Lindström, G.; Domingo, J. L. Perfluorinated chemicals in blood of residents in Catalonia (Spain) in relation to age and gender. *Environ. Int.* **2007**, *33*, 616–623.
- (60) Taniyasu, S.; Kannan, K.; So, K. M.; Gulkowska, A.; Sinclair, E.; Okazawa, T.; Yamashita, N. Analysis of fluorotelomer alcohols, fluorotelomer acids, and short- and long-chain

- perfluorinated acids in water and biota. *J. Chromatogr.* **2005**, *2093*, 89–97.
- (61) Falandysz, J.; Taniyasu, S.; Gulowska, A.; Yamashita, N.; Sculte-Oehlmann, U. Is fish a major source of fluorinated surfactants and repellents in humans living on the Baltic coast? *Environ. Sci. Technol.* **2006**, *40*, 748–751.
- (62) Kärman, A.; van Bavel, B.; Järnberg, U.; Hardell, L.; Lindström, G. Development of a solid-phase extraction-HPLC/single quadrupole MS method for quantification of perfluorochemicals in whole blood. *Anal. Chem.* **2005**, 864–870.
- (63) Kärman, A.; van Bavel, B.; Järnberg, U.; Hardell, L.; Lindström, G. Perfluorinated chemicals in relation to other persistent organic pollutants in human blood. *Chemosphere* **2006**, *64*, 1582–1591.
- (64) Li, X. M.; Yeung, L. W. Y.; Taniyasu, S.; Li, M.; Zhang, H.; Liu, D.; Lam, P. K. S.; Yamashita, N.; Dai, J. Perfluorooctanesulfonate and related fluorochemicals in the Amur tiger (*Panthera tigris altaica*) from China. *Environ. Sci. Technol.* **2008**, *42*, 7078–7083.
- (65) Van Leeuwen, S. P. J.; Kärman, A.; van Bavel, B.; De Boer, J.; Lindström, G. Struggle for quality in determination of perfluorinated contaminants in environmental and human samples. *Environ. Sci. Technol.* **2006**, *40*, 7854–7860.
- (66) Guruge, K. S.; Mange, P. M.; Yamanaka, N.; Miyazaki, S.; Taniyasu, S.; Yamashita, N. Species specific concentrations of perfluoroalkyl contaminants in farm and pet animals in Japan. *Chemosphere* **2008**, *73*, 210–215.
- (67) Ishibashi, H.; Iwata, H.; Kim, E.-Y.; Tao, L.; Kannan, K.; Amano, M.; Miyazaki, N.; Tanabe, S.; Valeriy, B. B.; Petrov, E. Contamination and effects of perfluorochemicals in Baikal seal (*Pusa sibirica*). 1. Residue level, tissue distribution and temporal trend. *Environ. Sci. Technol.* **2008**, *42*, 2295–2301.
- (68) Hart, K.; Kannan, K.; Isobe, T.; Takahashi, S.; Yamada, T. K.; Miyazaki, N.; Tanabe, S. Time trends and transplacental transfer of perfluorinated compounds in melon-headed whales stranded along the Japanese coast in 1982, 2001/2002 and 2006. *Environ. Sci. Technol.* **2008**, *42*, 7132–7137.
- (69) Furdui, V. I.; Helm, P. A.; Crozier, P. W.; Lucaciu, C.; Reiner, E. J.; Marvin, C. H.; Whittle, M. D.; Mabury, S. A.; Tomy, G. Temporal trends of perfluoroalkyl compounds with isomer analysis in lake trout from Lake Ontario (1974–2004). *Environ. Sci. Technol.* **2008**, *42*, 4739–4744.
- (70) Martin, J. W.; Mabury, S. A.; Solomon, K. R.; Muir, D. C. G. Bioconcentration and tissue distribution of perfluorinated acids in rainbow trouts (*Oncorhynchus mykiss*). *Environ. Toxicol. Chem.* **2003**, *22*, 196–204.
- (71) Conder, J. M.; Hoke, R. A.; De Wolf, W.; Russell, M. H.; Buck, R. C. Are PFCAs bioaccumulative? A critical review and comparison with regulatory lipophilic compounds. *Environ. Sci. Technol.* **2008**, *42*, 995–1003.
- (72) Keller, J.; Kannan, K.; Taniyasu, S.; Yamashita, N.; Day, R. D.; Arendt, M. D.; Segars, A. L.; Kucklick, J. R. Perfluorinated compounds in the plasma of loggerhead and Kemp's Ridley Sea turtles from the southeastern coast of the United States. *Environ. Sci. Technol.* **2005**, *39*, 9101–9108.
- (73) Kannan, K.; Perotta, E.; Thomas, N. J. Association between perfluorinated compounds and pathological conditions in Southern Sea otters. *Environ. Sci. Technol.* **2006**, *40*, 4943–4948.
- (74) Bossi, R.; Strand, J.; Sortkjær, O.; Larsen, M. M. Perfluoroalkyl compounds in Danish wastewater treatment plants and aquatic environments. *Environ. Int.* **2008**, *34*, 443–450.
- (75) Halldorsson, T. I.; Fei, C.; Olsen, J.; Lipworth, L.; McLaughlin, J. K.; Olsen, S. F. Dietary Predictors of Perfluorinated Chemicals: A Study from the Danish National Birth Cohort. *Environ. Sci. Technol.* **2008**, *42*, 8971–8977.
- (76) Spliethoff, H. M.; Tao, L.; Shaver, S. M.; Aldous, K. M.; Pass, K. A.; Kannan, K.; Eadon, G. A. Use of newborn screening program blood spots for exposure assessment: Declining levels of perfluorinated compounds in New York State infants. *Environ. Sci. Technol.* **2008**, *42*, 5361–5367.
- (77) Haug, L. S.; Thomsen, C.; Becher, G. Time trends and the influence of age and gender on serum concentrations of perfluorinated compounds in archived human samples. *Environ. Sci. Technol.* **2009**, *43* (6), 2131–2136.
- (78) Trudel, D.; Horowitz, L.; Wormuth, M.; Scheringer, M.; Cousins, I. T.; Hungerbühler, K. Estimating consumer exposure to PFOS and PFOA. *Risk Anal.* **2008**, *28*, 251–269.
- (79) Harada, K.; Saito, N.; Inoue, K.; Yoshinaga, T.; Watanabe, T.; Sasaki, S.; Kamiyama, S.; Koizumi, A. The influence of time, sex and geographic factors on levels of perfluorooctane sulfonate and perfluorooctanoate in human serum over the last 25 years. *J. Occup. Health* **2004**, *46*, 141–147.
- (80) Paul, A. G.; Jones, K. C.; Sweetman, A. J. A first global production, emission and environmental inventory for perfluorooctane sulfonate. *Environ. Sci. Technol.* **2008**, *43*, 386–392.
- (81) 3M Company. *3M Production History: Perfluorooctane sulfonyl fluoride (POSF)*; 2006; http://pops.int/documents/meetings/poprc/submissions/Comments_2006/3M.doc.
- (82) Toms, L. M.; Calafat, A. M.; Sjödin, A.; Harden, F.; Hobson, P.; Kato, K.; Thompson, J.; Mueller, J. F. Polyfluoroalkyl chemicals (PFCs) in Australian human blood serum. *Organohalogen Compd.* **2008**, *70*, 710–713.
- (83) Vestergren, R.; Cousins, I. T.; Trudel, D.; Wormuth, M.; Scheringer, M. Estimating the contribution of precursor compounds in consumer exposure to PFOS and PFOA. *Chemosphere* **2008**, *73*, 1617–1624.
- (84) D'Eon, J. C.; Mabury, S. A. Production of perfluorinated carboxylic acids (PFCAs) from the biotransformation of polyfluorinated phosphate surfactants (PAPS): Exploring the routes of human exposure. *Environ. Sci. Technol.* **2007**, *41* (13), 4799–4805.
- (85) Barber, J. L.; Berger, U.; Chaemfa, C.; Huber, S.; Jahnke, A.; Temme, C.; Jones, K. Analysis of perfluorinated alkyl substances in air samples from Northwest Europe. *J. Environ. Monit.* **2007**, 530–541.
- (86) Strynar, M. J.; Lindstrom, A. B. Perfluorinated compounds in house dust from Ohio and North Carolina, USA. *Environ. Sci. Technol.* **2008**, *42*, 3751–3756.
- (87) Fromme, H.; Schlummer, M.; Möller, A.; Gruber, L.; Wolz, G.; Ungewiss, J.; Böhmer, S.; Dekant, W.; Mayer, R.; Liebl, B.; Twardella, D. Exposure of an adult population to perfluorinated substances using duplicate diet portions and biomonitoring data. *Environ. Sci. Technol.* **2007**, *41*, 7928–7933.
- (88) Ericson, I.; Nadal, M.; van Bavel, B.; Lindström, G.; Domingo, J. L. Levels of perfluorochemicals in water samples from Catalonia, Spain: is drinking water a significant contribution to human exposure? *Environ. Sci. Pollut. R.* **2008**, *15*, 614–619.
- (89) Washburn, S. T.; Bingman, T. S.; Braithwaite, S. K.; Buck, R. C.; Buxton, W.; Clewell, H. J.; Haroun, L. A.; Kester, J.; Richard, R. W.; Shipp, A. Exposure assessment and risk characterization for perfluorooctanoate in selected consumer articles. *Environ. Sci. Technol.* **2005**, *39*, 3904–3910.
- (90) Saito, N.; Harada, K.; Inoue, K.; Sasaki, K.; Yoshinaga, T.; Koizumi, A. Perfluorooctane and perfluorooctane sulfonate concentrations in surface water in Japan. *J. Occup. Health* **2004**, *46*, 49–59.
- (91) Skutlarek, D.; Exner, M.; Färber, H. Perfluorinated Surfactants in Surface and Drinking Waters. *Environ. Sci. Pollut. R.* **2006**, *5*, 299–307.
- (92) Barton, C. A.; Butler, L. E.; Zarzecki, C. J.; Flaherty, J.; Kaiser, M. Characterizing perfluorooctanoate in ambient air near the fence line of a manufacturing facility: Comparing modeled and monitored values. *J. Air Waste Manage.* **2006**, *56*, 48–55.
- (93) Loos, R.; Wollgast, J.; Huber, T.; Hanke, G. Polar herbicides, pharmaceutical products, perfluorooctane sulfonate (PFOS), perfluorooctane acid (PFOA) and nonylphenol and its carboxylates and ethoxylates in surface and tap waters around Lake Maggiore in Northern Italy. *Anal. Bioanal. Chem.* **2007**, 1469–1478.
- (94) Plumlee, M. H.; Larabee, J.; Reinhard, M. Perfluorochemicals in water reuse. *Chemosphere* **2008**, *72*, 1541–1547.
- (95) Takagi, S.; Adachi, F.; Miyano, K.; Koizumi, Y.; Tanaka, H.; Mimura, M.; Watanabe, I.; Tanabe, S.; Kannan, K. Perfluorooctanesulfonate and perfluorooctanoate in raw and treated tap water from Osaka, Japan. *Chemosphere* **2008**, *72*, 1409–1412.
- (96) Begley, T. H.; White, K.; Honigfort, P.; Twaroski, M. L.; Neches, R.; Walker, R. A. Perfluorochemicals: Potential sources of and migration from food packaging. *Food Addit. Contam.* **2005**, *10*, 1023–1031.
- (97) Begley, T. H.; Hsu, W.; Noonan, G.; Diachenko, G. Migration of fluorochemical paper additives from food-contact paper into foods and food stimulants. *Food Addit. Contam.* **2007**, 1–7.
- (98) Shoeib, M.; Harner, T.; Lee, S. C.; Lane, D.; Zhu, J. Sorbent-impregnated polyurethane foam disk for passive air sampling of volatile fluorinated chemicals. *Anal. Chem.* **2008**, *80*, 675–682.
- (99) D'Eon, J. C.; Crozier, P. W.; Furdui, V. I.; Reiner, E. J.; Laurence Libelo, E.; Mabury, S. A. Observation of a commercial fluorinated material, the polyfluoroalkyl phosphoric acid diesters, in human sera, wastewater treatment plant sludge, and paper fibers. *Environ. Sci. Technol.* **2009**, *43* (12), 4589–4594.

- (100) Tittlemier, S. A.; Pepper, K.; Seymour, C.; Moisey, J.; Bronsson, R.; Cao, X. L.; Dabeka, R. W. Dietary exposure of Canadians to perfluorinated carboxylates and perfluorooctane sulfonate via consumption of meat, fish and fast food items prepared in their packaging. *J. Agric. Food Chem.* **2007**, *55*, 3203–3210.
- (101) Del Gobbo, L.; Tittlemier, S. A.; Diamond, M.; Pepper, K.; Tague, B.; Yeudall, F.; Vanderlinden, L. Cooking decreases observed perfluorinated compound concentrations in fish. *J. Agric. Food Chem.* **2008**, *56*, 7551–7559.
- (102) Berger, U.; Holmström, K.; Glynn, A.; Berglund, M.; Ankarberg, E.; Törnkvist, A.; Perfluorierade alkylsyror i matkorsprover och fisk från Vättern och Ålands hav. Rapport till Naturvårdsverket, Programområde Miljögiftssamordning. Dnr: 721-5953-06Mm.
- (103) Gulkowska, A.; Jiang, Q.; So, K. M.; Taniyasu, H.; Lam, P. K. S.; Yamashita, N. Persistent Perfluorinated Acids in Seafood Collected from Two Cities of China. *Environ. Sci. Technol.* **2006**, *40*, 3736–3741.
- (104) Kärman, A.; Harada, K. H.; Inoue, K.; Takasuga, T.; Etumasa, O.; Koizumi, A. Relationship between dietary exposure and serum perfluorochemical (PFC) levels-A case study. *Environ. Int.* **2009**, *35*, 712–717.
- (105) Gruber, L.; Schlummer, M.; Ungewiss, J.; Wolz, G.; Moeller, A.; Weise, N.; Fromme, H. Analysis of sub-ppb levels of perfluorooctanoic acid (PFOA) and perfluorooctanesulfonate (PFOS) in food and fish. *Organohalogen Compd.* **2007**, *69*, 142–145.
- (106) Stahl, T.; Heyn, J.; Thiele, H.; Hütter, J.; Failing, K.; Georgii, S.; Brunn, H. Carryover of Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonate (PFOS) from Soil to Plants. *Arch. Environ. Contam. Toxicol.* **2008**, DOI 10.1007/s00244-008-9272-9.
- (107) Renner, R. EPA finds record PFOS, PFOA levels in Alabama grazing fields. *Environ. Sci. Technol.* **2009**, *5*, 1245–1246.
- (108) Kubwabo, C.; Stewart, B.; Zhu, J. P.; Marro, L. Occurrence of perfluorosulfonates and other perfluorochemicals in dust from selected homes in the city of Ottawa, Canada. *J. Environ. Monit.* **2005**, *7*, 1074–1078.
- (109) Moriwaki, H. Y.; Takatah, Y.; Arakawa, R. Concentrations of Perfluorooctane sulfonate (PFOS) and Perfluorooctanoic acid (PFOA) in vacuum cleaner dust collected in Japanese homes. *J. Environ. Monit.* **2003**, 753–757.
- (110) Björklund, J.; Thuresson, K.; de Wit, C. A. Perfluoroalkyl compounds (PFCs): Concentrations, human exposure estimates and sources. *Environ. Sci. Technol.* **2009**, *42*, 2277–2281.
- (111) Fromme, H.; Nitschke, L.; Kiranoglu, M.; Albrecht, M.; Völkel, W. Perfluorinated substances in house dust in Bavaria, Germany. *Organohalogen Compd.* **2008**, *70*, 1048–1050.
- (112) Boulanger, B.; Vargo, J. D.; Schnoor, J. L.; Hornbuckle, K. Evaluation of perfluorooctane surfactants in a wastewater treatment system and in a commercial surface protection product. *Environ. Sci. Technol.* **2005**, *39*, 5524–5530.
- (113) Tittlemier, S. A.; Pepper, K.; Edwards, L. Concentrations of perfluorooctanesulfonamides in Canadian total diet study composite food samples collected between 1992 and 2004. *J. Agric. Food Chem.* **2006**, *54*, 8385–8389.
- (114) Kemper, R. A. *Perfluorooctanoic acid: Toxicokinetics in rat*; Report DuPont-7473; Association of Plastics Manufacturers of Europe; Brussels, Belgium, 2003.
- (115) Kennedy, G. L., Jr.; Hall, C. T.; Brittelli, M. R.; Barnes, J. R.; Chen, H. C. Inhalation toxicity of ammoniumperfluorooctanoate. *Food Chem. Toxicol.* **1986**, *24*, 1325–1329.
- (116) Hinderliter, P. M.; DeLorme, M. P.; Kennedy, G. L. Perfluorooctanoic acid: Relationship between repeated inhalation exposures and plasma PFOA concentration in the rat. *Toxicology* **2006**, *222*, 80–85.
- (117) Fasano, W. J.; Kennedy, G. L.; Szostek, B.; Farrar, D. G.; Ward, R. J.; Haroun, L.; Hinderliter, P. M. Penetration of ammonium perfluorooctanoate through rat and human skin in vitro. *Drug Chem. Toxicol.* **2005**, *28*, 79–90.
- (118) Hanhijärvi, H.; Ylinen, M.; Kojo, A.; Kosma, V. Elimination and toxicity of perfluorooctanoic acid during subchronic administration in the Wistar rat. *Pharmacol. Toxicol.* **1987**, *61*, 66–68.
- (119) Ylinen, M.; Kojo, A.; Hanhijärvi, H.; Peura, P. Disposition of perfluorooctanoic acid in the rat after single dose and subchronic administration. *Bull. Environ. Contam. Toxicol.* **1990**, *44*, 45–53.
- (120) Hanhijärvi, H. A proposed species difference in the renal excretion of perfluorooctanoic acid in the beagle dog and rat. In *New Developments in Biosciences: Their Implications for Laboratory Animal Science*; Beynen, A. C., Solleveld, H. A., Eds.; Martinus Nijhoff: Dordrecht, 1988; pp 404–412.
- (121) Kudo, N.; Suzuki, E.; Katakura, M.; Ohmori, K.; Noshiro, R.; Kawashima, Y. Comparison of the elimination between perfluorinated fatty acids with different chain length in rats. *Chem. Biol. Interact.* **2001**, *134*, 203–216.
- (122) Van den Heuvel, J. P.; Kuslikis, B. I.; Van Rafelghem, M. J.; Peterson, M. J. Tissue distribution, metabolism and elimination of perfluorooctanoic acid in male and female rats. *J. Biochem. Toxicol.* **1991**, *6*, 83–92.
- (123) Ohmori, K.; Kudo, N.; Katayama, K.; Kawashima, Y. Comparison of the toxicokinetics between perfluorocarboxylic acids with different carbon chain length. *Toxicology* **2003**, *184*, 135–140.
- (124) Butenhoff, J. L.; Kennedy, H.; Hinderliter, P. M.; Lieder, P. H.; Jung, R.; Hansen, K. J.; Gorman, G. S.; Noker, P. E.; Thomford, P. J. Pharmacokinetics in cynomolgus monkeys. *Toxicol. Sci.* **2004**, *83*, 394–406.
- (125) Ward, K. V.; Erhardt, P.; Bachman, K. Application of simple mathematic expressions to relate the half-lives of xenobiotics in rats to humans. *J. Pharmacol. Toxicol.* **2005**, *51*, 57–64.
- (126) Ward, K. W.; Smith, B. R. A comprehensive quantitative and qualitative evaluation of extrapolation of intravenous pharmacokinetic pharmacokinetic parameters from rat, dog and monkey to humans I. Clearance. *Drug. Metab. Dispos.* **2004**, *32*, 603–611.
- (127) Andersen, M. E.; Clewell, H. J., III; Tan, Y.-M.; Butenhoff, J. L.; Olsen, G. W. Pharmacokinetic modeling of saturable, renal resorption perfluoroalkylacids in monkeys-Probing the determinants of long plasma half-lives. *Toxicology* **2006**, *227*, 156–164.
- (128) Cahill, T. M.; Cousins, I. T.; MacKay, D. Development of a generalized physiologically based pharmacokinetic model for multiple environmental contaminants. *Environ. Toxicol. Chem.* **2003**, *22*, 26–34.
- (129) Harada, K.; Inoue, K.; Morikawa, A.; Yoshinaga, T.; Saito, N.; Koizumi, A. Renal clearance of perfluorooctane sulfonate and perfluorooctanoate and their species-specific excretion. *Environ. Res.* **2005**, *99*, 253–261.
- (130) PERFluorinated Organics in Our Diet (PERFOOD). A three-year collaborative project, starting early 2009. Financed by the Food, Agriculture and Fisheries Programme of the EU Seventh Framework Programme (SF7) (Workprogramme: FP7-KBBE-2007-2B, Workprogramme Topic: KBBE-2008-2-4-01). Coordinator: Dr. Pim de Voogt, University of Amsterdam, NL.
- (131) Telomer Research Program, 2002. Telomer research program update; Presented at the USEPA OPPT, November 25, 2002; US Environmental Protection Agency Public Docket AR226-1141.
- (132) Armitage, J.; Cousins, I. T.; Buck, R. C.; Prevedouros, K.; Russel, M. H.; Macleod, M.; Korzeniowski, S. H. Modeling global-scale fate and transport of perfluorooctanoate emitted from direct sources. *Environ. Sci. Technol.* **2006**, *40*, 6969–6975.

ES900228K