

## Spatial and Temporal Patterns in Concentrations of Perfluorinated Compounds in Bald Eagle Nestlings in the Upper Midwestern United States

William T. Route,<sup>\*,†</sup> Robin E. Russell,<sup>‡</sup> Andrew B. Lindstrom,<sup>§</sup> Mark J. Strynar,<sup>§</sup> and Rebecca L. Key<sup>†</sup>


<sup>†</sup>U.S. National Park Service, Great Lakes Inventory and Monitoring Network, Suite D, 2800 Lake Shore Drive East, Ashland, Wisconsin 54806, United States

<sup>‡</sup>U.S. Geological Survey, National Wildlife Health Center, 6006 Schroeder Road, Madison, Wisconsin 53711, United States

<sup>§</sup>National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina 27711, United States

### Supporting Information

**ABSTRACT:** Perfluorinated chemicals (PFCs) are of concern due to their widespread use, persistence in the environment, tendency to accumulate in animal tissues, and growing evidence of toxicity. Between 2006 and 2011 we collected blood plasma from 261 bald eagle nestlings in six study areas from the upper Midwestern United States. Samples were assessed for levels of 16 different PFCs. We used regression analysis in a Bayesian framework to evaluate spatial and temporal trends for these analytes. We found levels as high as 7370 ng/mL for the sum of all 16 PFCs ( $\Sigma$ PFCs). Perfluorooctanesulfonate (PFOS) and perfluorodecanesulfonate (PFDS) were the most abundant analytes, making up 67% and 23% of the PFC burden, respectively. Levels of  $\Sigma$ PFC, PFOS, and PFDS were highest in more urban and industrial areas, moderate on Lake Superior, and low on the remote upper St. Croix River watershed. We found evidence of declines in  $\Sigma$ PFCs and seven analytes, including PFOS, PFDS, and perfluorooctanoic acid (PFOA); no trend in two analytes; and increases in two analytes. We argue that PFDS, a long-chained PFC with potential for high bioaccumulation and toxicity, should be considered for future animal and human studies.



### INTRODUCTION

Perfluorinated compounds (PFCs) have been in worldwide production and use since the 1950s. They have the unique properties of repelling both water and oil making them useful in a variety of products, including the production of polymers for nonstick coatings in cookware and food packaging, water and stain repellants for textiles, and in the manufacture of lubricants, firefighting foams, and pesticide formulations. The first reports of their presence in human blood were published in the late 1960s and 1970s<sup>1,2</sup> with indications of significant environmental contamination by the late 1970s.<sup>3</sup> At this same time laboratory studies of perfluorooctanesulfonate (PFOS) and perfluorooctanoic acid (PFOA) in primates indicated a wide range of deleterious effects including hepatotoxicity, cancer, and death at relatively low dose rates.<sup>4</sup> Industry researchers began to express concern by the late 1970s<sup>5,6</sup> leading to studies on the health effects in occupationally exposed workers.<sup>7</sup> PFCs are now ubiquitous across the globe, and many studies have documented their persistence, rates of accumulation in animals, and effects on human health,<sup>8</sup> including child behavior.<sup>9</sup> Scientists, regulators, and managers are therefore in need of information on the spatial patterns and temporal trends in PFCs.

Monitoring PFC concentrations in wildlife has proven helpful in estimating distributions,<sup>10</sup> long-term trends,<sup>11</sup> and routes of human exposure.<sup>12,13</sup> In particular, sentinel species such as the bald eagle (*Haliaeetus leucocephalus*), which feeds high on the trophic food web, are excellent sentinels for PFC contamination.<sup>14,15</sup> Eagles provide several tissues (e.g., blood, feathers, and eggs) that are relatively easy to obtain<sup>16,17</sup> and nestlings are particularly useful for monitoring local pollution.<sup>17</sup> Garrett et al.<sup>18</sup> reviewed several studies and estimated home range size of nesting bald eagles varied from 1.5 km<sup>2</sup> to 21.7 km<sup>2</sup>. Hence contaminants found in their young are indicative of contamination within 2.6 km of the nesting site.


The bald eagle ranged across most of North America but declined in the late 1950s and 1960s due to indiscriminant killing, habitat loss, and chemical contamination. Their populations increased after DDT (dichlorodiphenyltrichloroethane) was banned in the U.S. in 1972 and Canada in 1989. They were removed from the U.S. List of Endangered and

Received: March 4, 2014

Revised: May 5, 2014

Accepted: May 20, 2014

Published: June 5, 2014



**Figure 1.** Location of six study areas in the upper Midwestern United States where bald eagle nestlings were sampled for perfluorinated compounds, 2006–2011.

Threatened Wildlife in 2007.<sup>19</sup> Concern remains for this species, however, due to its vulnerability to persistent pollutants. In this study we report on PFC concentrations in blood plasma of bald eagle nestlings from the upper Midwest. Our objectives were to evaluate the spatial patterns and temporal trends of 16 PFC analytes in this region and to suggest implications of exposure to humans and wildlife.

## MATERIALS AND METHODS

**Field Collections.** From 2006 to 2011 we collected blood samples from occupied bald eagle nests in six study areas: the Apostle Islands National Lakeshore (APIS); Wisconsin's Lake Superior South Shore (LSSS); the upper St. Croix National Scenic Riverway (U-SACN); the lower St. Croix National Scenic Riverway (L-SACN); the Mississippi National River and Recreation Area (MISS); and a portion of Pools 3 and 4 of the Mississippi River (Pools 3 & 4) (Figure 1). From mid-May through early June each year, when nestlings were five to nine weeks old, we climbed to occupied nests, hand-captured the nestlings, and brought them to the ground for sampling. Nestlings were weighed, aged, banded, sampled, and placed back in the nest. All measurements were consistent with other investigators in the upper Midwest.<sup>20–22</sup> Nestling age was determined by the length of the eighth primary feather,<sup>23</sup> and sex was determined by PCR-based genetic analysis.<sup>24</sup>

We took  $\leq 11$  mL of blood from the brachial vein of nestlings using a sterile, 10 mL polypropylene syringe, and 20-gauge needle. The blood sample was immediately transferred to a sterile 10 mL vacutainer. Within 12 h of collection, samples were centrifuged at 1200 rpm for 10–12 min to separate

plasma from red blood cells. A sterile glass pipet was used to transfer  $\leq 1.0$  mL of plasma to a polypropylene vial for analysis. Glass pipettes were previously baked at 650 °F (343 °C) to remove chemical residues. A sample of syringes, needles, vacutainers, and vials were tested by the 3M Environmental Laboratory, Maplewood, MN and verified free of PFCs. Plasma samples were kept frozen until delivered to an analytical laboratory.

All nestlings (1–4) were sampled from each nest unless they were too young or old to handle. A single sample was chosen from each nest for analyses in this study (arbitrarily in 2006, randomly thereafter). The remaining samples were archived frozen ( $-20$  °C) for future analyses.

The Wisconsin State Laboratory of Hygiene (WSLH) in Madison, WI was the primary analytical laboratory. To measure interlaboratory variability we conducted blind studies with the U.S. Environmental Protection Agency (USEPA) laboratory in Research Triangle Park, NC and the 3M Environmental Laboratory in Maplewood, MN. Each laboratory worked independently, was unaware of results from other laboratories, and did not know the location of sampled nestlings.

**Laboratory Procedures.** Each laboratory used high performance liquid chromatography/tandem mass spectrometry and gradient elution chromatography to measure concentrations of up to 16 PFCs, including the following: perfluorobutanoic acid (PFBA), perfluorobutanesulfonate (PFBS), perfluorodecanoic acid (PFDA), perfluorododecanoic acid (PFDoA), perfluorodecanesulfonate (PFDS), perfluoroheptanesulfonate (PFHpS), perfluoroheptanoic acid (PFHpA), perfluorohexanoic acid (PFHxA), perfluorohexanesulfonate

(PFHxS), perfluorononanoic acid (PFNA), perfluorooctanoic acid (PFOA), perfluorooctanesulfonate (PFOS), perfluorotetradecanoic acid (PFTeDA), perfluoropentanoic acid (PFPA), perfluorotridecanoic acid (PFTTrDA), and perfluoroundecanoic acid (PFuDA). The 16 analytes were selected because they were measurable using standard laboratory procedures and believed to be present in the region.

Each laboratory used slightly different procedures and had different limits of quantification (LOQ) (see Supporting Information (SI) Table S11). All three laboratories had previous experience with PFC analyses and used matrix spikes of known concentrations to assess accuracy and surrogate spikes to evaluate extraction efficiency. Results from each laboratory are known only to the study authors.

**Statistical Analyses.** Twelve of the 16 PFCs were detected by two or more laboratories at levels above their respective LOQ in  $\geq 59\%$  of their samples, and these were used for all analyses. Four analytes (PFBS, PFHpA, PFHxA, and PFPA) were included in the  $\sum$ PFC but were omitted from further analyses because they were detected in  $<25\%$  of the samples. These four contributed  $<1\%$  of the total PFC burden.

We estimated the spatial distribution and temporal changes in PFCs using regression analysis in a Bayesian statistical framework<sup>25,26</sup> to account for variability in laboratories, missing values, potential differences due to age and sex, and for measurements that were below a laboratory's LOQ. We accounted for the observed PFC levels from the three laboratories by formulating log-transformed PFC levels ( $Y$ ) for each individual nest ( $i$ ) and each laboratory ( $l$ ) as a multivariate normal distribution. The multivariate normal distribution is formulated in WinBUGS<sup>27</sup> as  $Y_{i,l} \sim \text{MVN}(\mu_{i,l}, \Omega)$ ; where  $\Omega$  is the precision matrix for the vector of random components, i.e. the values from each of the three laboratories. The prior for  $\Omega$  is a Wishart distribution. For values below LOQ we bounded the upper limit of  $\mu_{i,l}$  by the lowest LOQ from the laboratory.

PFC levels were modeled as a function of a spatial correlation term, a time effect, a time by space effect, an effect of eagle sex, an effect of eagle age, and a fixed effect of laboratory (weight was highly correlated with age so was not included). Priors for the estimates of covariate coefficients were based on a noninformative normal distribution  $\beta \sim \mathcal{N}(0, 0.001)$ . To account for spatial correlation between eagle territories we used the correlated autoregressive model, *car.normal*, in WinBUGS.<sup>28</sup> We considered all Lake Superior nests, and the nearest upstream and downstream nests on rivers, as "neighbors" with spatial weights of one; all other nests were weighted zero. The time by territory effect was formulated as a random variable  $\mathcal{N}(\mu, \tau)$  where  $\mu$  is the mean value of the random effect, and  $\tau$  is the precision parameter  $1/\sigma$ . We ran three chains for 50000 iterations and discarded the first 25000 as burn-in. Every 10th value in the chains was used to estimate posterior distributions of the estimated parameter coefficients. We assessed convergence using the Gelman-Rubin statistic and R-hat values using library "coda".<sup>29</sup> We conducted a Goodness of Fit analysis on estimates from the hierarchical model against residuals of observed PFOS values from all laboratories as a check on model sensitivity ( $R^2 = 0.81$ ).

The Bayesian framework allowed us to interpret the credible intervals (CI) as true probabilities of change in levels of an analyte. We concluded there was "strong evidence of change" if  $\geq 90\%$  of the posterior estimates were above or below zero, "moderate evidence of change" if  $>80\%$  but  $<90\%$  of the

estimates were above or below zero, "weak" evidence if  $>70\%$  but  $<80\%$  were above or below zero, and "no evidence" of change if  $<70\%$  were above or below zero. An analyte was determined to be either increasing or decreasing if  $>70\%$  of the posterior estimates were above or below zero, respectively.

## RESULTS AND DISCUSSION

**Sample Collection.** From 2006 through 2011 we collected blood plasma from 261 bald eagle nestlings in six study areas (Table 1). The number of samples for each area varied yearly

**Table 1. Distribution Across Study Areas and Years for Bald Eagle Nestlings Sampled and Analyzed for PFC Analytes**

study area <sup>a</sup>	2006 <sup>b</sup>	2007 <sup>b</sup>	2008 <sup>b</sup>	2009 <sup>c</sup>	2010	2011	totals
APIS	8	6	5	0	9	9	37
LSSS	0	6	4	0	0	1	11
U-SACN	11	8	0	0	11	12	32
L-SACN	3	4	7	9	13	13	49
MISS	11	11	15	18	23	20	98
Pools 3 & 4	0	0	15	12	4	2	34
totals	33	35	46	39	51	57	261

<sup>a</sup>APIS = Apostle Islands National Lakeshore, LSSS = Lake Superior South Shore, U-SACN = upper St. Croix National Scenic Riverway, L-SACN = lower St. Croix National Scenic Riverway, MISS = Mississippi National River and Recreation Area, Pools 3 & 4 = portions of pools 3 and 4 on the Mississippi River. <sup>b</sup>Two way blind laboratory analyses; all samples analyzed by laboratories 1 and 2. <sup>c</sup>Three way blind laboratory analyses; all samples analyzed by laboratories 1, 2, and 3.

due to size of the eagle population, nest occupancy, and funding. All samples were analyzed by the primary laboratory, samples from 2006 through 2008 were analyzed by two laboratories ( $n = 114$ ), and samples from 2009 were analyzed by three laboratories ( $n = 39$ ).


**Effects of Laboratory and Nestling Age.** For each of the 12 analytes we calculated correlation coefficients (CC) between paired laboratories, the intraclass correlation coefficient (ICC) between all three laboratories, and intralab coefficients of variation (CV) (Table S12). Those analytes found in high concentrations were reproducible and consistent between laboratories. PFOS and PFDS made up  $>90\%$  of the PFC burden, and they, along with the sum of all 16 PFCs, hereafter  $\sum$ PFC, had CCs and ICCs  $>0.80$  and within-laboratory CVs  $<26$ . The less abundant analytes varied more widely with CCs and ICCs ranging from  $-0.12$  to  $0.97$  and CVs of  $12.33$  to  $71.55$ . These results are similar to other interlaboratory comparisons where large differences have been found for analytes found at extremely low concentrations.<sup>30</sup> However, most CCs in our study (68%) were above 0.80 indicating that, while some laboratory measurements differed in magnitude, they trended in the same direction.

We found strong evidence (the 95% CI did not include zero) that levels of three analytes were affected by nestling age: the estimated coefficient for age for PFNA was 1.17 (95% CI = 1.02–1.35) and for PFUnA was 1.23 (95% CI = 1.00–1.27) indicating an increase with age for these two analytes while the coefficient for PFBA was 0.77 (95% CI = 0.60–1.00) indicating a decline with age in this analyte. Other PFCs appeared unaffected by age. The mechanisms involved in age-related differences likely include the bioaccumulative properties of each analyte, exposure time, and the development of excretory

**Table 2. Estimated Geometric Mean and Range (ng/mL) in Concentrations of 12 PFCs in Blood Plasma of Bald Eagle Nestlings**

analyte	geometric mean and (range) PFC concn (ng/mL) <sup>a</sup>					
	APIS	LSSS	U-SACN	L-SACN	MISS	Pools 3 & 4
∑PFC <sup>b</sup>	552 (139–1420)	490 (109–472)	163 (14.5–205)	644 (60.8–3450)	607 (62.2–7370)	941 (579–2930)
PFOS	265 (71.0–830)	425 (75.5–290)	77.5 (6.56–180)	429 (10.0–2400)	421 (45.0–4200)	800 (414–1400)
PFDS	13.6 (LOQ–100)	13.7 (0.65–7.80)	3.95 (LOQ–20.0)	131 (13.0–1090)	79.8 (1.90–4100)	265 (130–1400)
PFDA	12.6 (0.06–77.0)	11.3 (3.92–29.0)	8.49 (1.10–5.19)	11.3 (2.95–30.0)	11.3 (2.20–85.0)	12.3 (LOQ–37.0)
PFUnA	17.5 (20.4–110)	11.8 (10.9–81.9)	7.79 (1.30–10.4)	9.46 (2.17–19.0)	10.44 (1.70–49.6)	9.55 (2.30–65.0)
PFDoA	7.03 (3.54–27.0)	5.81 (1.98–16.2)	3.18 (0.04–1.90)	5.47 (1.10–18.0)	6.10 (0.90–33.0)	5.54 (2.11–31.0)
PFNA	8.13 (21.0–160)	4.93 (5.57–83.0)	4.15 (0.65–8.39)	4.31 (0.29–12.0)	4.62 (LOQ–19.0)	4.18 (LOQ–11.0)
PFTrDA	3.87 (9.0–63.0)	3.01 (4.40–48.0)	2.18 (0.13–5.80)	2.68 (0.64–14.0)	2.73 (0.48–14.0)	2.59 (0.94–12.0)
PFHpS	1.88 (0.58–5.40)	1.97 (0.48–1.80)	1.11 (LOQ–2.90)	1.97 (0.19–4.40)	1.96 (0.23–16.0)	2.60 (1.76–11.0)
PFHxS	2.25 (LOQ–8.60)	1.80 (LOQ–8.55)	2.73 (LOQ–9.10)	1.81 (LOQ–8.30)	1.43 (LOQ–46.7)	0.77 (LOQ–26.0)
PFTeDA	1.49 (0.84–19.0)	1.42 (0.43–16.0)	1.22 (LOQ–2.40)	1.43 (0.24–14.0)	1.41 (0.28–310)	1.95 (LOQ–14.0)
PFOA	1.01 (LOQ–14.0)	0.64 (LOQ–5.30)	0.37 (LOQ–1.49)	0.34 (LOQ–10.0)	0.52 (LOQ–9.90)	0.49 (LOQ–14.6)
PFBA	0.31 (LOQ–22.0)	0.43 (LOQ–0.78)	0.50 (LOQ–0.90)	0.47 (LOQ–32.0)	0.55 (LOQ–78.0)	0.47 (LOQ–5.60)

<sup>a</sup>Means are modeled from independent measurements by three laboratories; min and max values are the actual highest and lowest value measured by any one laboratory. <sup>b</sup>∑PFC is the sum of all 16 PFC analytes measured by the independent laboratories; four analytes that made up <1% of the sample volume and were below LOQ >25% of the time were excluded from summary statistics.



**Figure 2.** Geometric mean concentrations of ∑PFCs in bald eagle nestlings, 2006–2011. Values are modeled from measurements by three independent laboratories. Categories were selected using natural breaks in Arc GIS. A = location of 3M PFC production facility; B = Minneapolis/St. Paul WWTP and St. Paul Downtown Airport; C = the communities of St. Croix Falls, WI and Taylors Falls, MN. See Figure 1 for study area acronyms.

organs and processes in the growing nestling. We found no effect on PFC levels due to nestling sex.

**PFC Concentrations and Spatial Patterns.** The geometric mean (GM) of ∑PFCs was highest at Pools 3 & 4 (941 ng/mL) followed by L-SACN (644 ng/mL), MISS (607 ng/mL), APIS (552 ng/mL), and LSSS (490 ng/mL), with the

lowest concentrations occurring on the remote U-SACN (163 ng/mL) (Table 2). This pattern of higher ∑PFC concentrations near urban centers has been found in a range of environmental matrices.<sup>31–33</sup> In a state-wide assessment of PFCs in Minnesota fish, five PFCs were found at higher levels near Minneapolis/St. Paul, MN than in more rural areas of the

state.<sup>33</sup> Similarly, 10 PFCs in surface water of the Cape Fear River Basin, NC were found at higher levels near known or suspected sources in urban areas,<sup>31</sup> and in East Asia, studies have shown higher levels of PFCs in air above industrialized regions.<sup>34</sup>

There are several sources of PFCs in this region<sup>35</sup> but most notable is the 3M Cottage Grove facility (Figure 2, A) which used electrochemical fluorination (ECF) to make perfluorooctane sulfonyl fluoride (PSOF) until 2002.<sup>36</sup> PSOF was the starting material for production of PFOS and other PFC analytes. The ECF process is relatively unrefined, and other PFCs may have been unintentional byproducts. Contamination around this<sup>37</sup> and other PFC production facilities has been well documented in soil, water, sediments, and fish.<sup>14,38,39</sup> Two bald eagle territories downriver of this 3M facility (at 8.6 km and 13.8 km) had the highest mean concentrations of  $\sum$ PFCs over the six year study. Moreover, in 2011, one nestling in this area of river had 7370 ng/mL  $\sum$ PFC, the highest level we are aware of in bald eagles.

Concentrations of PFCs in nestlings were high along the Mississippi River even upstream from the 3M facility, however. Compared to upstream samples, we observed a near doubling of  $\sum$ PFC concentrations in nestlings beginning near the Minneapolis/St. Paul wastewater treatment plant (WWTP) where treated effluent is discharged to the river (Figure 2, B). Lee et al.<sup>40</sup> found high levels of other organic compounds downstream from this same WWTP where metal plating industries, known for using PFCs, contribute to the influent. Also in this area is the St. Paul Downtown Airport, which is within the flood plain of the river. Other investigators have documented PFCs in surface water near airports where they are used in fire-fighting foams.<sup>41</sup> Determining the source of PFCs to the river is further complicated, however, by the presence of landfills where 3M disposed of PFC waste since the 1950s. Four groundwater aquifers below these landfills are known to be contaminated with PFCs<sup>42</sup> and may release them to the river.

We found a similar pattern on the St. Croix River where  $\sum$ PFCs increased sharply immediately downstream from the communities of Taylors Falls and St. Croix Falls (Figure 2, C). This section of river is subjected to effluent from several WWTPs serving communities and industry along the lower St. Croix valley.

The APIS and LSSS study areas are comparatively remote, yet nestlings there had moderately high  $\sum$ PFC levels. We suspect this is due in part to Lake Superior's physical characteristics. Lake Superior has a 31700 mi<sup>2</sup> (82103 km<sup>2</sup>) surface area that absorbs airborne contaminants from global and regional sources, and, though sparsely populated, there are numerous WWTPs from municipalities along its 2700 mile (4385 km) shoreline with >200 tributaries that serve many communities. Moreover, the 191 year residence time for water in Lake Superior results in ample time for bioconcentration (resuspension of contaminated sediments and recycling through the food web). Furthermore, bald eagles on Lake Superior have been shown to feed on gulls and other piscivorous birds at higher rates than inland eagles, which further biomagnifies contaminants.<sup>43</sup> This slow removal and bioaccumulation of persistent contaminants from Lake Superior's food web has been demonstrated with DDE (dichlorodiphenyldichloroethylene) and PCBs (polychlorinated biphenyl), which remain at relatively high levels in bald

eagle nestlings more than three decades after being banned in North America.<sup>44</sup>

**Patterns of Individual Analytes.** The 95% credible intervals (in Bayesian statistics these are analogous to traditional confidence intervals) overlap for all PFCs in all study areas (Table S13). Nonetheless, five general patterns emerge (Table 2): (1) PFOS was found in all sampled nestlings and was the most abundant PFC (GM = 77.5–800 ng/mL) in all study areas, contributing 67% of the total PFC burden; (2) PFDS was the second most abundant PFC (GM = 3.95–265 ng/mL), accounting for 23% of burden, and highest in the urbanized, riverine study areas (L-SACN, MISS, and Pools 3 & 4 were more than 5-fold higher than APIS and LSSS and 20-fold higher than U-SACN); (3) PFOA was generally at low concentrations (GM = 0.34–1.01 ng/mL) but highest in Lake Superior study areas (APIS and LSSS); (4) The Lake Superior study areas had higher levels of more analytes than other study areas (highest or second highest for 9 of 12 PFCs); and (5) The U-SACN had the lowest mean concentrations of nearly all PFCs (9 of 12).

We found PFOS at the highest mean concentrations at Pools 3 & 4 (GM = 800 ng/mL), though the highest recorded value for an individual nestling was at MISS (4200 ng/mL) and the second highest at L-SACN (2400 ng/mL). Geometric means for MISS, L-SACN, and LSSS were similar (421–429 ng/mL), APIS was moderately high at 265 ng/mL, and the remote U-SACN study area had the lowest concentrations (77.5 ng/mL). The only comparable study on bald eagle nestlings in the region<sup>45</sup> reported measurable PFOS plasma concentrations in 32 of 33 nestlings from the Great Lakes between 1990 and 1993 with an arithmetic mean of 330 ng/mL (SE = 126). However, the spike recovery of PFOS was only 17% in that study, suggesting concentrations may have been much higher. Nonetheless, this and the current investigation make it clear that eagles on the Great Lakes have had high burdens of PFOS at least since 1990. Moreover, PFOS is the predominant PFC found throughout the aquatic food web in the upper Midwest. Elevated levels have been found in water, benthic organisms, fish, turtles, mink, and tissue from moribund bald eagles from the Great Lakes.<sup>46,47</sup> Similar to our findings, high levels have been reported in water, sediments, invertebrates, and fish from the Mississippi River below Minneapolis and St. Paul, MN with the highest concentrations occurring below the 3M Cottage Grove facility.<sup>33,37,48,49</sup>

Newsted<sup>50</sup> calculated a toxicity reference value (TRV) of 1700 ng PFOS/mL blood-plasma as protective of a level IV fish-eating bird such as an eagle irrespective of sex and reproductive status. We found GM concentrations to be below this TRV in all of our study areas; however, levels for some individual nestlings were higher: 5 of 98 (5.1%) at MISS and 2 of 21 (9.5%) at L-SACN. We found no effects of PFOS on bald eagle productivity in our study. We did not measure potential sublethal pathological, physiological, or behavioral effects at the individual level.

The second most abundant PFC was PFDS, accounting for 23% of the total burden. PFDS was highest in nestlings along the Mississippi and lower St. Croix Rivers (Pools 3 & 4, MISS, and L-SACN; GM = 80–265 ng/mL), moderate in nestlings from Lake Superior (APIS and LSSS; GM = 13.7–17.5 ng/mL), and lowest on the upper St. Croix River drainage (U-SACN; GM = 3.95) (Table 2). The high PFDS levels we found in the urban study areas are potentially significant. Few studies report on PFDS in environmental or human samples. Furdul et

al.<sup>47</sup> found PFDS in 89% of lake trout from the Great Lakes and found levels to be correlated with PFOS - both being highest in Lake Erie and lowest in Lake Superior lake trout. However, the National Health and Nutritional Examination Study, which tests a representative sample of 5000 people across the U.S. biannually, does not include PFDS.<sup>51</sup> Given that longer-chain PFCs like PFDS tend to be more toxic and prone to bioaccumulation,<sup>52</sup> we argue that it should be considered in future animal and human studies.

We found PFOA concentrations to be comparatively low overall, and the highest levels were in nestlings from the Lake Superior study areas (Table 2). The low levels in bald eagles are likely due to PFOA's low bioaccumulation properties,<sup>46</sup> and the higher levels in Lake Superior nestlings may be due to differences in availability. PFOA is generally found at higher levels than PFOS in surface waters of the Great Lakes,<sup>46</sup> including Lake Superior where Scott et al. found PFOA to be highest among 23 analytes measured in surface water.<sup>53</sup> The authors of this latter study estimated that 35% of the PFOA in surface water was from precipitation and 59% was from tributaries, noting that WWTPs, many which are located on tributaries, concentrated PFOA up to 20-fold that of intake water. By comparison, median concentrations of PFOA was fourth among 13 PFCs tested in surface water of the upper Mississippi River.<sup>48</sup>

Other major analytes included PFDA (GM = 8.49–12.6 ng/mL), PFUnA (7.79–17.5 ng/mL), PFNA (4.15–8.13 ng/mL), and PFDoA (3.18–7.03 ng/mL). All other PFCs were found in concentrations <3.87 ng/mL.

**Temporal Trends.** Over the six year study we found strong evidence of decline in  $\sum$ PFCs and five analytes (*probability*  $\geq$  90% to 100%), moderate evidence of decline for two analytes (*probability*  $\geq$  80% to <90%), no evidence of change in three analytes (*probability* > 30% to <70%), and evidence of increase in two analytes (*probability of a decline* = 0 and the expected ratio of change >1.10; Table 3). However, these trends were not uniform across study areas (Table SI4). For example, the probability that PFHpS declined from 2006 to 2011 was >92% for all study areas except Pools 3 & 4 where evidence of decline

was lacking (*probability* = 53%). Conversely, evidence of PFOS decline was strong at Pools 3 & 4 (*probability* = 95%), moderate at MISS and L-SACN (*probability* = 83% to 89%), weak at APIS and LSSS (*probability* = 73% to 76%), and lacking at U-SACN (*probability of decline* = 47%). Similar declines in PFOS have been documented by others in water, sediments, and fish along the MISS section of the Mississippi River between 2004 and 2012.<sup>37</sup> These declines are likely the result of 3M's discontinuation of PFOS production in 2002. Nonetheless, this food web will continue to contain PFOS and other PFCs for many years. This is due to the extent of local soil, sediment, and groundwater contamination, leaching from PFC-containing products in municipal landfills, and the continued production of traditional and alternative PFCs globally. Even in the U.S. a limited amount of restricted PFC production continues where there are no known alternatives.<sup>54</sup> Moreover, resuspension of PFCs by flooding can make them available to the aquatic food web. For example, in 2011, even after strong evidence of declines, we found the highest concentration of  $\sum$ PFCs in a nestling (7370 ng/mL). This followed a spring of flooding on the Mississippi River which was higher and lasted longer (well into the bald eagle nesting period) compared to the prior 30 years (unpublished USGS streamflow data at Anoka, MN).

Trends for PFNA and PFDA also differed among study areas. We found moderate evidence of declines (*probability* = 80% to 90%) for PFNA at Pools 3 & 4 and U-SACN, no evidence of decline at MISS and LSSS, (*probability* < 70%), and evidence of increase at APIS (*probability of increase* = 83%). Similarly, evidence of decline in PFDA was lacking in all study areas except at U-SACN where it showed evidence of increase (*probability of increase* 83%).

We detected strong evidence that PFTeDA and PFTrDA were increasing in nestlings from all study areas (*probability of increase* > 90%). The expected ratios of change were >1.0, and the mean expected ratio was 2.5 for PFTeDA indicating that, on average, levels of this contaminant were more than doubling every year (Table 3). These two PFCs made up <0.5% of the total PFC load in nestlings; however, the increasing levels and lack of knowledge about their source and toxicity suggests a need for increased study.

In summary, this six year study documents extensive contamination of aquatic systems of the upper Midwest by at least 16 different PFC analytes. Bald eagle nestlings served as good biosentinels of local contamination, and our results further substantiate findings of others that PFC levels in the environment are linked to effluent from municipal wastewater systems and industrial waste. Bayesian modeling provided a robust means of including measurements from independent laboratories and allowed us to estimate the probabilities of increase or decline for each of 12 PFCs at six study areas. While we found that many PFCs declined during this study, their persistence and the continued production of new alternative PFCs suggest the need for continued research.

## ■ ASSOCIATED CONTENT

### § Supporting Information

We provide four tables with information on laboratory methods, comparisons among laboratories, midranges and confidence intervals, and probabilities of declines at each study area. This material is available free of charge via the Internet at <http://pubs.acs.org>.

**Table 3. Expected Ratio of Change and the 95% Credible Interval (C.I.) for 12 PFC Analytes Found in Bald Eagle Nestlings**

analyte <sup>a</sup>	expected ratio of change (95% C.I.)	probability levels are declining <sup>b</sup>
PFOA	0.78 (0.72, 0.86)	100
PFUnA	0.85 (0.77–0.93)	100
PFHxS	0.83 (0.73–0.95)	100
PFDS	0.83 (0.69–1.01)	97
PFDoA	0.91 (0.82–1.01)	96
$\sum$ PFCs	0.92(0.82–1.03)	90
PFHpS*	0.78 (0.57–1.43)	86
PFOS	0.95 (0.86–1.06)	82
PFNA	0.97 (0.87–1.09)	64
PFBA*	1.01 (0.83–1.24)	42
PFDA	1.02 (0.92–1.14)	33
PFTeDA*	2.48 (1.42–5.03)	0
PFTrDA*	1.30 (1.10–1.51)	0

<sup>a</sup>An asterisk (\*) indicates analytes with four years (2008–2011) of data rather than 6;  $n = 202$  nestlings. <sup>b</sup>The probability that levels are declining, calculated as the percentage of trend estimates from the posterior distribution that are below zero.

## AUTHOR INFORMATION

### Corresponding Author

\*Phone: 715-682-0631 ext 221. E-mail: bill\_route@nps.gov.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

Primary funding was provided by the U.S. National Park Service Great Lakes Inventory and Monitoring Network with additional support from the Minnesota Pollution Control Agency, the Great Lakes Restoration Initiative, and the Donald Weesner Foundation. A portion of this work was conducted as part of a Cooperative Research and Development Agreement between the USEPA and Agilent Technologies (#437-07). P. Bost, USEPA, and the 3M Corporation provided blind laboratory analyses, and G. Miller, Prairie Island Indian Community provided sample results and field assistance, as in-kind contributions. The Wisconsin State Laboratory of Hygiene was contracted as the primary analytical laboratory. We thank J. Campbell-Spickler and G. Renzullo for tree climbing and handling of nestlings, M. Martell for handling of nestlings, and many national park service staff and volunteers for assistance with data collection and logistics. We appreciate constructive comments on drafts of this manuscript by B. La Francois, D. VanderMeulen, L. Weller, and R. Erikson. The United States Environmental Protection Agency through its Office of Research and Development has reviewed this article and approved it for publication. Any use of trade, product, or firm names are for descriptive purposes only and do not imply endorsement by the U.S. Government.

## REFERENCES

- Taves, D. Evidence that there are two forms of fluoride in human serum. *Nature* **1968**, *217* (133), 1050–1051.
- Taves, D. R.; Grey, W. S.; Brey, W. S. Organic fluoride in human plasma - its distribution and partial identification. *Toxicol. Appl. Pharmacol.* **1976**, *37* (1), 120–120.
- Dupont Memorandum: C-8 Environmental Status Administrative Record AR226-2000. U.S. Environmental Protection Agency: 1983.
- Goldenthal, E. I.; Jessup, D. C.; Geil, R. G.; Mehring, J. S. *Ninety-day subacute rhesus monkey toxicity study. Fluorad Fluorochemical Surfactant FC-95. Study No. 137-092*. International Research and Development Corporation: Mattawan, MI, U.S. EPA Docket No. AR226-0137, 1978.
- Dupont Meeting with 3M May 30. US EPA Administrative Record, AR226-1449. 1978.
- Belisle, J. Organic fluorine in human serum - natural versus industrial sources. *Science* **1981**, *212* (4502), 1509–1510.
- Steenland, K.; Fletcher, T.; Savitz, D. A. Epidemiologic evidence on the health effects of perfluorooctanoic acid (PFOA). *Environ. Health Perspect.* **2010**, *118* (8), 1100–1108.
- Post, G. B.; Cohn, P. D.; Cooper, K. R. Perfluorooctanoic acid (PFOA), an emerging drinking water contaminant: A critical review of recent literature. *Environ. Res.* **2012**, *116*, 93–117.
- Gump, B. B.; Wu, Q.; Dumas, A. K.; Kannan, K. Perfluorochemical (PFC) exposure in children: associations with impaired response inhibition. *Environ. Sci. Technol.* **2011**, *45* (19), 8151–8159.
- Houde, M.; De Silva, A. O.; Muir, D. C. G.; Letcher, R. J. Monitoring of perfluorinated compounds in aquatic biota: an updated review. *Environ. Sci. Technol.* **2011**, *45* (19), 7962–7973.
- Rotander, A.; Karrman, A.; Van Bavel, B.; Polder, A.; Riget, F.; Auousson, G. A.; Vikingsson, G.; Gabrielsen, G. W.; Bloch, D.; Dam, M. Increasing levels of long-chain perfluorocarboxylic acids (PFCAs) in Arctic and North Atlantic marine mammals, 1984–2009. *Chemosphere* **2012**, *86* (3), 278–285.
- Holzer, J.; Midasch, O.; Rauchfuss, K.; Kraft, M.; Reupert, R.; Angerer, J.; Kleeschulte, P.; Marschall, N.; Wilhelm, M. Biomonitoring of perfluorinated compounds in children and adults exposed to perfluorooctanoate-contaminated drinking water. *Environ. Health Perspect.* **2008**, *116* (5), 651–657.
- Falandysz, J.; Taniyasu, S.; Gulkowska, A.; Yamashita, N.; Schulte-Oehlmann, U. Is fish a major source of fluorinated surfactants and repellents in humans living on the Baltic Coast? *Environ. Sci. Technol.* **2006**, *40*, 748–751.
- Kannan, K.; Franson, J. C.; Bowerman, W. W.; Hansen, K. J.; Jones, P. D.; Giesy, J. P. Perfluorooctane sulfonate in fish-eating water birds including bald eagles and albatrosses. *Environ. Sci. Technol.* **2001**, *35* (15), 3065–3070.
- Dauwe, T.; Van De Vijver, K.; De Coen, W.; Eens, M. PFOS levels in the blood and liver of a small insectivorous songbird near a fluorochemical plant. *Environ. Int.* **2006**, *33* (3), 357–361.
- Gebbink, W. A.; Letcher, R. J.; Hebert, C. E.; Weseloh, D. V. C. Twenty years of temporal change in perfluoroalkyl sulfonate and carboxylate contaminants in herring gull eggs from the Laurentian Great Lakes. *J. Environ. Monit.* **2011**, *13* (12), 3365–3372.
- Holmstrom, K. E.; Johansson, A.-K.; Bignert, A.; Lindberg, P.; Berger, U. Temporal trends of perfluorinated surfactants in Swedish peregrine falcon eggs (*Falco peregrinus*), 1974–2007. *Environ. Sci. Technol.* **2010**, *44* (11), 4083–4088.
- Garrett, M. G.; Watson, J. W.; Anthony, R. G. Bald eagle home range and habitat use in the Columbia River Estuary. *J. Wildl. Manage.* **1993**, *57* (1), 19–27.
- Removing the bald eagle in the lower 48 states from the List of Endangered and Threatened Wildlife and Plants. *Fed. Regist.* **2007**, *72* (130), 37346–37371.
- Bowerman, W. W. Regulation of bald eagle productivity in the Great Lakes Basin: an ecological and toxicological approach. Ph.D. Dissertation, Michigan State University, East Lansing, MI, 1993.
- Bowerman, W. W.; Best, D. A.; Giesy, J. P.; Shieldcastle, M. C.; Meyer, M. W.; Postupalsky, S. J.; Sikarskie, J. G. Associations between regional differences in polychlorinated biphenyls and dichlorodiphenyldichloroethylene in blood of nestling bald eagles and reproductive productivity. *Environ. Toxicol. Chem.* **2003**, *22* (2), 371–376.
- Dykstra, C. R.; Meyer, M. W.; Rasmussen, P. W.; Warnke, D. K. Contaminant concentrations and reproductive rate of Lake Superior bald eagles, 1989–2001. *J. Great Lakes Res.* **2005**, *31* (2), 227–235.
- Bortolotti, G. R. Criteria for determining age and sex of nestling bald eagles. *J. Field Ornithol.* **1984**, *55* (4), 467–481.
- Boutette, J. B.; Ramsay, E. C.; Potgieter, L. N.; Kania, S. A. An improved polymerase chain reaction-restriction fragment length polymorphism assay for gender identification in birds. *J. Avian Med. Surg.* **2002**, *16* (3), 198–202.
- Bayes, T. An essay toward solving a problem in the doctrine of chances. *Philos. Trans. R. Soc. London* **1764**, *53*, 370–418.
- Gelman, A.; Carlin, J. B.; Stern, H. S.; Rubin, D. B. *Bayesian data analysis*, 2nd ed.; Chapman and Hall/CRC: Boca Raton, FL, 2003.
- Lunn, D. J.; Thomas, A.; Best, N.; Spiegelhalter, D. WinBUGS-a Bayesian modelling framework: concepts, structure, and extensibility. *Stat. Comput.* **2000**, *10* (4), 325–337.
- Besag, J.; York, J.; Mollie, A. Bayesian image restoration with two applications in spatial statistics. *Ann. Inst. Stat. Math.* **1991**, *43*, 1–59.
- Plummer, M.; Best, N.; Cowles, K.; Vines, K. CODA: Convergence diagnosis and output analysis for MCMC. *R News* **2006**, *6* (1), 7–11.
- Longnecker, M. P.; Smith, C. S.; Kissling, G. E.; Hoppin, J. A.; Butenhoff, J. L.; Decker, E.; Ehresman, D. J.; Ellefson, M. E.; Flaherty, J.; Gardner, M. S.; Langlois, E.; Leblanc, A.; Lindstrom, A. B.; Reagen, W. K.; Strynar, M. J.; Studabaker, W. B. An interlaboratory study of perfluorinated alkyl compound levels in human plasma. *Environ. Res.* **2008**, *107* (2), 152–159.

- (31) Nakayama, S.; Strynar, M. J.; Helfant, L.; Egeghy, P.; Ye, X.; Lindstrom, A. B. Perfluorinated compounds in the Cape Fear Drainage basin in North Carolina. *Environ. Sci. Technol.* **2007**, *41* (15), 5271–5276.
- (32) Zushi, Y.; Ye, F.; Motegi, M.; Nojiri, K.; Hosono, S.; Suzuki, T.; Kosugi, Y.; Yaguchi, K.; Masunaga, S. Spatially detailed survey on pollution by multiple perfluorinated compounds in the Tokyo Bay basin of Japan. *Environ. Sci. Technol.* **2011**, *45* (7), 2887–2893.
- (33) Delinsky, A. D.; Mccann, P. J.; Varns, J. L.; Mcmillan, L.; Nakayama, S. F.; Lindstrom, A. B. Geographical distribution of perfluorinated compounds in fish from Minnesota lakes and rivers. *Environ. Sci. Technol.* **2010**, *44* (7), 2549–2554.
- (34) Saito, T.; Yokouchi, Y.; Stohl, A.; Taguchi, S.; Mukai, H. Large emissions of perfluorocarbons in East Asia deduced from continuous atmospheric measurements. *Environ. Sci. Technol.* **2010**, *44* (11), 4089–4095.
- (35) Ye, X.; Schoenfuss, H. L.; Jahns, N. D.; Delinsky, A. D.; Strynar, M. J.; Varns, J.; Nakayama, S. F.; Helfant, L.; Lindstrom, A. B. Perfluorinated compounds in common carp (*Cyprinus carpio*) filets from the Upper Mississippi River. *Environ. Int.* **2008**, *34* (7), 932–8.
- (36) Monson, B. A.; Solem, L.; Hoff, P.; Hora, M.; Mccann, P.; Briggs, M.; Stiras, J.; Delain, S. *Mississippi River Pool 2 intensive study of perfluorochemicals and water: 2009*; Minnesota Pollution Control Agency, St. Paul, MN, 2010.
- (37) Monson, B. A. *Perfluorochemicals in Mississippi River Pool 2:2012 update*; Minnesota Pollution Control Agency, St. Paul, MN, 2013.
- (38) Emmett, E. A.; Shofer, F. S.; Zhang, H.; Freeman, D.; Desai, C.; Shaw, L. M. Community exposure to perfluorooctanoate: relationships between serum concentrations and exposure sources. *J. Occup. Environ. Med.* **2006**, *48* (8), 759–770.
- (39) Hansen, K. J.; Johnson, H. O.; Eldridge, J. S.; Butenhoff, J. L.; Dick, L. A. Quantitative characterization of trace levels of PFOS and PFOA in the Tennessee River. *Environ. Sci. Technol.* **2002**, *36*, 1681–1685.
- (40) Lee, K. E.; Barber, L. B.; Furlong, E. T.; Cahill, J. D.; Kolpin, D. W.; Meyer, M. W.; Zaugg, S. D. *Presence and distribution of organic wastewater compounds in wastewater, surface, ground, and drinking waters, Minnesota, 2000-02*; Minnesota Department of Health and Minnesota Pollution Control Agency, St. Paul, MN, 2004; pp 1–33.
- (41) Saito, N.; Sasaki, K.; Nakatome, K.; Harada, K.; Yoshinaga, T.; Koizumi, A. Perfluorooctane sulfonate concentrations in surface water in Japan. *Arch. Environ. Contam. Toxicol.* **2003**, *45* (2), 149–158.
- (42) *Public health assessment: perfluorochemical contamination in southern Washington County, northern Dakota County, and southeastern Ramsey County, Minnesota*; Minnesota Department of Health, St. Paul, MN, 2012; p 161.
- (43) Kozie, K. D.; Anderson, R. K. Productivity, diet, and environmental contaminants in bald eagles nesting near the Wisconsin shoreline of Lake Superior. *Arch. Environ. Contam. Toxicol.* **1991**, *20* (1), 41–48.
- (44) Dykstra, C. R.; Route, W. T.; Meyer, M. W.; Rasmussen, P. W. Contaminant concentrations in bald eagles nesting on Lake Superior, the upper Mississippi River, and the St. Croix River. *J. Great Lakes Res.* **2010**, *36* (3), 561–569.
- (45) Kannan, K.; Franson, J. C.; Bowerman, W. W.; Hansen, K. J.; Jones, P. D.; Giesy, J. P. Perfluorooctane Sulfonate in Fish-Eating Water Birds Including Bald Eagles and Albatrosses. *Environ. Sci. Technol.* **2001**, *35* (15), 3065–3070.
- (46) Giesy, J.; Mabury, S.; Martin, J.; Kannan, K.; Jones, P.; Newsted, J.; Coady, K. Perfluorinated compounds in the Great Lakes. In *Persistent organic pollutants in the Great Lakes*; Hites, R. A., Ed.; Springer-Verlag: Berlin, Heidelberg, Germany, 2006; pp 391–438.
- (47) Furdui, V. I.; Stock, N. L.; Ellis, D. A.; Butt, C. M.; Whittle, D. M.; Crozier, P. W.; Reiner, E. J.; Muir, D. C.; Maybury, S. A. Spatial distribution of perfluoroalkyl contaminants in lake trout from the Great Lakes. *Environ. Sci. Technol.* **2007**, *41* (5), 1554–1559.
- (48) Nakayama, S. F.; Strynar, M. J.; Reiner, J. L.; Delinsky, A. D.; Lindstrom, A. B. Determination of perfluorinated compounds in the Upper Mississippi River Basin. *Environ. Sci. Technol.* **2010**, *44*, 4103–4109.
- (49) Ye, X.; Strynar, M. J.; Nakayama, S. F.; Varns, J.; Helfant, L.; Lazorchak, J.; Lindstrom, A. B. Perfluorinated compounds in whole fish homogenates from the Ohio, Missouri, and Upper Mississippi Rivers, USA. *Environ. Pollut.* **2008**, *156* (3), 1227–1232.
- (50) Newsted, J. L.; Jones, P. D.; Coady, K.; Giesy, J. P. Avian toxicity reference values for perfluorooctane sulfonate. *Environ. Sci. Technol.* **2005**, *39* (23), 9357–9362.
- (51) Kato, K.; Wong, L.-Y.; Jia, L. T.; Kuklenyik, Z.; Calafat, A. M. Trends in exposure to polyfluoroalkyl chemicals in the U.S. population: 1999–2008. *Environ. Sci. Technol.* **2011**, *45* (19), 8037–8045.
- (52) Conder, J. M.; Hoke, R. A.; Wolf, W. D.; Russell, M. H.; Buck, R. C. Are PFCA's bioaccumulative? A critical review and comparison with regulatory criteria and persistent lipophilic compounds. *Environ. Sci. Technol.* **2008**, *42* (4), 995–1003.
- (53) Scott, B. F.; De Silva, A. O.; Spencer, C.; Lopez, E.; Backus, S. M.; Muir, D. C. G. Perfluoroalkyl acids in Lake Superior water: trends and sources. *J. Great Lakes Res.* **2010**, *36* (2), 277–284.
- (54) Oliaei, F.; Kriens, D.; Weber, R.; Watson, A. PFOS and PFC releases and associated pollution from a PFC production plant in Minnesota (USA). *Environ. Sci. Pollut. Res.* **2013**, *20* (4), 1977–1992.

#### NOTE ADDED AFTER ISSUE PUBLICATION

This paper was published June 5, 2014. In the version published July 9, 2014, the display of authors and the addresses have been revised.